

**bs-9223R**

**[ Primary Antibody ]**

**Bioss**  
ANTIBODIES

www.bioss.com.cn

sales@bioss.com.cn

techsupport@bioss.com.cn


400-901-9800

## RNF70 Rabbit pAb


### DATASHEET

<b>Host:</b> Rabbit	<b>Isotype:</b> IgG	<b>Applications:</b> <b>WB</b> (1:500-2000) <b>IHC-P</b> (1:100-500) <b>IHC-F</b> (1:100-500) <b>IF</b> (1:50-200)
<b>Clonality:</b> Polyclonal		
<b>GeneID:</b> 64219	<b>SWISS:</b> Q8NG27	
<b>Target:</b> RNF70		
<b>Immunogen:</b> KLH conjugated synthetic peptide derived from human RNF70: 541-640/643.		
<b>Purification:</b> affinity purified by Protein A		
<b>Concentration:</b> 1mg/ml		<b>Reactivity:</b> Human, Rat (predicted: Mouse, Rabbit, Pig, Dog, Horse)
<b>Storage:</b> 0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol. Shipped at 4°C. Store at -20°C for one year. Avoid repeated freeze/thaw cycles.		<b>Predicted MW.:</b> 71 kDa
<b>Background:</b> Has E2-dependent E3 ubiquitin-protein ligase activity. Ubiquitinates MAGED1 antigen leading to its subsequent degradation by proteasome (By similarity). May be involved in protein sorting.		<b>Subcellular Location:</b> Cytoplasm ,Nucleus

### VALIDATION IMAGES



Sample: 293T(Human) Cell Lysate at 30 ug  
HepG2(Human) Cell Lysate at 30 ug Primary:  
Anti-RNF70 (bs-9223R) at 1/1000 dilution  
Secondary: IRDye800CW Goat Anti-Rabbit IgG at  
1/20000 dilution Predicted band size: 71 kD  
Observed band size: 71 kD



Paraformaldehyde-fixed, paraffin embedded (Rat brain); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Antibody incubation with (RNF70) Polyclonal Antibody, Unconjugated (bs-9223R) at 1:400 overnight at 4°C, followed by operating according to SP Kit(Rabbit) (sp-0023) instructions and DAB staining.