## bs-9223R

## [ Primary Antibody ]

www.bioss.com.cn sales@bioss.com.cn techsupport@bioss.com.cn

DATASHEET -

Host: Rabbit Isotype: IgG

Clonality: Polyclonal

RNF70 Rabbit pAb

GenelD: 64219 SWISS: Q8NG27

Target: RNF70

**Immunogen:** KLH conjugated synthetic peptide derived from human RNF70:

541-640/643.

**Purification:** affinity purified by Protein A

Concentration: 1mg/ml

**Storage:** 0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50%

Glycerol.

Shipped at 4°C. Store at -20°C for one year. Avoid repeated

freeze/thaw cycles.

Background: Has E2-dependent E3 ubiquitin-protein ligase activity.

Ubiquitinates MAGED1 antigen leading to its subsequent degradation by proteasome (By similarity). May be involved in

protein sorting.

Applications: WB (1:500-2000)

400-901-9800

**IHC-P** (1:100-500) **IHC-F** (1:100-500) **IF** (1:100-500)

Reactivity: Human, Rat

(predicted: Mouse, Rabbit,


Pig, Dog, Horse)

**Predicted** 


71 kDa MW.:

**Subcellular Location:** Cytoplasm ,Nucleus

## VALIDATION IMAGES



Sample: 293T(Human) Cell Lysate at 30 ug HepG2(Human) Cell Lysate at 30 ug Primary: Anti-RNF70 (bs-9223R) at 1/1000 dilution Secondary: IRDye800CW Goat Anti-Rabbit IgG at 1/20000 dilution Predicted band size: 71 kD Observed band size: 71 kD



Paraformaldehyde-fixed, paraffin embedded (Rat brain); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Antibody incubation with (RNF70) Polyclonal Antibody, Unconjugated (bs-9223R) at 1:400 overnight at 4°C, followed by operating according to SP Kit(Rabbit) (sp-0023) instructions and DAB staining.