

**bs-17654R****[ Primary Antibody ]****SPERT Rabbit pAb****BioSS**  
**ANTIBODIES**

www.bioss.com.cn



sales@bioss.com.cn

techsupport@bioss.com.cn

400-901-9800

**— DATASHEET —****Host:** Rabbit**Isotype:** IgG**Clonality:** Polyclonal**GeneID:** 220082**SWISS:** Q8NA61**Target:** SPERT**Immunogen:** KLH conjugated synthetic peptide derived from human SPERT: 1-100/448.**Purification:** affinity purified by Protein A**Concentration:** 1mg/ml**Storage:** 0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol.

Shipped at 4°C. Store at -20°C for one year. Avoid repeated freeze/thaw cycles.

**Background:** A variety of morphological and molecular changes are required for spermatozoa formation. These steps are temporally guided by the transcription and translation of several testis-specific genes. SPERT (spermatid associated), also known as CBY2 (chibby homolog 2), spermatid flower-like structure protein or NURIT, is a 448 amino acid novel leucine-zipper protein belonging to the chibby family of proteins. Expressed uniquely in the spermatid flower-like structure, SPERT interacts with Nek1, a member of the NIMA-family kinase family that is associated centrosomal stability and ciliogenesis. Containing a leucine-zipper motif and two coiled-coil regions, SPERT is transcribed through the elongation stage of the spermatids. SPERT is absent from mature spermatozoa and is thought to be involved in transporting proteins that are to be discarded via the residual bodies.**Applications:** **WB** (1:500-2000)**IHC-P** (1:100-500)**IHC-F** (1:100-500)**IF** (1:100-500)**Reactivity:** Human, Rat  
(predicted: Mouse, Rabbit, Sheep, Cow, Horse)**Predicted**  
**MW.:** 52 kDa**Subcellular**  
**Location:** Cytoplasm**— VALIDATION IMAGES —**

Sample: HeLa(human) Cell Lysate at 40 ug  
DU145(human) Cell Lysate at 40 ug  
Primary: Anti-SPERT(bs-17654R) at 1/300 dilution  
Secondary: IRDye800CW Goat Anti-Rabbit IgG at 1/20000 dilution  
Predicted band size: 52kD  
Observed band size: 52kD

Paraformaldehyde-fixed, paraffin embedded (rat kidney tissue); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Antibody incubation with (SPERT) Polyclonal Antibody, Unconjugated (bs-17654R) at 1:400 overnight at 4°C, followed by a conjugated secondary (sp-0023) for 20 minutes and DAB staining.