

bs-17662R**[Primary Antibody]****SPHKAP Rabbit pAb****Bioss**
ANTIBODIES

www.bioss.com.cn

sales@bioss.com.cn

techsupport@bioss.com.cn

400-901-9800

— DATASHEET —

Host: Rabbit	Isotype: IgG	Applications: IHC-P (1:100-500) IHC-F (1:100-500) IF (1:100-500) ICC/IF (1:100-500) ELISA (1:5000-10000) Reactivity: (predicted: Human, Mouse, Rat, Rabbit, Pig, Sheep, Cow, Zebrafish, Chicken, Dog, Horse) Predicted MW.: 186 kDa Subcellular Location: Cytoplasm
Clonality: Polyclonal		
GeneID: 80309	SWISS: Q2M3C7	
Target: SPHKAP		
Immunogen: KLH conjugated synthetic peptide derived from human SPHKAP: 1131-1230/1700.		
Purification: affinity purified by Protein A		
Concentration: 1mg/ml		
Storage: 0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol. Shipped at 4°C. Store at -20°C for one year. Avoid repeated freeze/thaw cycles.		
Background: Anchoring protein that binds preferentially to the type I regulatory subunit of c-AMP-dependent protein kinase (PKA type I) and targets it to distinct subcellular compartments. May act as a converging factor linking cAMP and sphingosine signaling pathways. Plays a regulatory role in the modulation of SPHK1.		

— SELECTED CITATIONS —

- **[IF=4.183]** Zhu Y et al. Protein Expression Profile in Rat Silicosis Model Reveals Upregulation of PTPN2 and Its Inhibitory Effect on Epithelial-Mesenchymal Transition by Dephosphorylation of STAT3. Int J Mol Sci. 2020 Feb 11;21(4). WB ;Rat&Mouse. 32054021