



## Angiotensin II type 1A receptor Rabbit pAb

Catalog Number: bs-2132R

Target Protein: Angiotensin II type 1A receptor

Concentration: 1mg/ml

Form: Liquid Host: Rabbit

Clonality: Polyclonal

Isotype: IgG

Applications: WB (1:500-2000), IHC-P (1:100-500), IHC-F (1:100-500), IF (1:100-500)

Reactivity: Human, Mouse, Rat

Predicted MW: 41 kDa Entrez Gene: 185 Swiss Prot: P30556

Source: KLH conjugated synthetic peptide derived from human Angiotensin II type 1A receptor:

101-200/359.

Purification: affinity purified by Protein A

Storage: 0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol.

Shipped at 4°C. Store at -20°C for one year. Avoid repeated freeze/thaw cycles.

Background: Angiotensin II is a potent vasopressor hormone and a primary regulator of aldosterone

secretion. It is an important effector controlling blood pressure and volume in the

cardiovascular system. It acts through at least two types of receptors. This gene encodes the type 1 receptor which is thought to mediate the major cardiovascular effects of angiotensin

II. This gene may play a role in the generation of reperfusion arrhythmias following restoration of blood flow to ischemic or infarcted myocardium. It was previously thought


only one type 1 receptor gene in humans. At least five transcript variants have been

that a related gene, denoted as AGTR1B, existed; however, it is now believed that there is


described for this gene. Additional variants have been described but their full-length nature has not been determined. The entire coding sequence is contained in the terminal exon and

is present in all transcript variants. [provided by RefSeq].


## **VALIDATION IMAGES**




Sample: Lane 1: Liver (Mouse) Lysate at 40 ug Lane 2: Heart (Mouse) Lysate at 40 ug Lane 3: Placenta (Mouse) Lysate at 40 ug Lane 4: Kidney (Mouse) Lysate at 40 ug Lane 5: Embryo (Mouse) Lysate at 40 ug Lane 6: Cerebrum (Mouse) Lysate at 40 ug Lane 7: NIH/3T3 (Mouse) cell Lysate at 30 ug Lane 8: Liver (Rat) Lysate at 40 ug Lane 9: Heart (Rat) Lysate at 40 ug Lane 10: Kidney (Rat) Lysate at 40 ug Lane 11: Cerebrum (Rat) Lysate at 40 ug Lane 12: Embryo (Rat) Lysate at 40 ug Primary: Anti-Angiotensin II type 1A receptor (bs-2132R) at 1/1000 dilution Secondary: IRDye800CW Goat Anti-Rabbit IgG at 1/20000 dilution Predicted band size: 43 kD Observed band size: 46 kD




Sample: A549(Human) Cell Lysate at 430 ug Primary: Anti-Angiotensin II type 1A receptor (bs-2132R) at 1/500 dilution Secondary: IRDye800CW Goat Anti-Rabbit IgG at 1/20000 dilution Predicted band size: 41 kD Observed band size: 41/51 kD



Sample: Liver (Mouse) Lysate at 40 ug Lung (Mouse) Lysate at 40 ug Adrenal gland (Rat) Lysate at 40 ug Primary: Anti-Angiotensin II type 1A receptor (bs-2132R) at 1/1000 dilution Secondary: IRDye800CW Goat Anti-Rabbit IgG at 1/20000 dilution Predicted band size: 41 kD Observed band size: 56 kD



Paraformaldehyde-fixed, paraffin embedded (Rat brain); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Antibody incubation with (Angiotensin II type 1A receptor) Polyclonal Antibody, Unconjugated (bs-2132R) at 1:400 overnight at 4°C, followed by operating according to SP Kit(Rabbit) (sp-0023) instructions and DAB staining.



Paraformaldehyde-fixed, paraffin embedded (rat kidney); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Antibody incubation with (Angiotensin II type 1A receptor) Polyclonal Antibody, Unconjugated (bs-2132R) at 1:200 overnight at 4°C, followed by operating according to SP Kit(Rabbit) (sp-0023) instructions and DAB staining.

## PRODUCT SPECIFIC PUBLICATIONS

[IF=2.03] Liu, Junjun, et al. "Differential effects of angiotensin II receptor blockers on Aβ generation." Neuroscience Letters (2014). WB; ="Mouse" .24680748