## bs-2132R

# [ Primary Antibody ]

www.bioss.com.cn sales@bioss.com.cn techsupport@bioss.com.cn 400-901-9800

# Angiotensin II type 1A receptor Rabbit pAb

DATASHEET -

Host: Rabbit Isotype: IgG

Clonality: Polyclonal

GenelD: 185 **SWISS:** P30556

Target: Angiotensin II type 1A receptor

Immunogen: KLH conjugated synthetic peptide derived from human

Angiotensin II type 1A receptor: 101-200/359. < Extracellular >

Purification: affinity purified by Protein A

Concentration: 1mg/ml

Storage: 0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50%

Glycerol.

Shipped at 4°C. Store at -20°C for one year. Avoid repeated

freeze/thaw cycles.

**Background:** Angiotensin II is a potent vasopressor hormone and a primary regulator of aldosterone secretion. It is an important effector controlling blood pressure and volume in the cardiovascular system. It acts through at least two types of receptors. This gene encodes the type 1 receptor which is thought to mediate the major cardiovascular effects of angiotensin II. This gene may play a role in the generation of reperfusion arrhythmias following restoration of blood flow to ischemic or infarcted myocardium. It was previously thought that a related gene, denoted as AGTR1B, existed; however, it is now believed that there is only one type 1 receptor gene in humans. At least five transcript variants have been described for this gene. Additional variants have been described but their fulllength nature has not been determined. The entire coding sequence is contained in the terminal exon and is present in all transcript variants. [provided by RefSeq].

Applications: WB (1:500-2000)


**IHC-P** (1:100-500) **IHC-F** (1:100-500) **IF** (1:100-500)

Reactivity: Human, Mouse, Rat

Predicted 41 kDa MW.:

**Subcellular** Cell membrane


### VALIDATION IMAGES -




Sample: Liver (Mouse) Lysate at 40 ug Lung (Mouse) Lysate at 40 ug Adrenal gland (Rat) Lysate at 40 ug Primary: Anti-Angiotensin II type 1A receptor (bs-2132R) at 1/1000 dilution Secondary: IRDye800CW Goat Anti-Rabbit IgG at 1/20000 dilution Predicted band size: 41 kD Observed band size: 56 kD



Sample: A549(Human) Cell Lysate at 40 ug Primary: Anti-Angiotensin II type 1A receptor (bs-2132R) at 1/500 dilution Secondary: IRDye800CW Goat Anti-Rabbit IgG at 1/20000 dilution Predicted band size: 41 kD Observed band size: 41/51 kD



Paraformaldehyde-fixed, paraffin embedded (rat kidney); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Antibody incubation with (Angiotensin II type 1A receptor) Polyclonal Antibody, Unconjugated (bs-2132R) at 1:200 overnight at 4°C, followed by operating according to SP Kit(Rabbit) (sp-0023) instructions and DAB staining.



Paraformaldehyde-fixed, paraffin embedded (Rat brain); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Antibody incubation with (Angiotensin II type 1A receptor) Polyclonal Antibody, Unconjugated (bs-2132R) at 1:400 overnight at 4°C, followed by operating according to SP Kit(Rabbit) (sp-0023) instructionsand DAB staining.

# - SELECTED CITATIONS -

• [IF=2.03] Liu, Junjun, et al. "Differential effects of angiotensin II receptor blockers on Aβ generation." Neuroscience Letters (2014). WB; Mouse. 24680748