bs-23231R

[Primary Antibody]

ALK Rabbit pAb



www.bioss.com.cn sales@bioss.com.cn techsupport@bioss.com.cn 400-901-9800

– DATASHEET –––––		400-901-9800
Host: Rabbit	Isotype: IgG	Applications: WB (1:500-2000)
Clonality: Polyclonal		Beactivity: Mouse (predicted: Human
GenelD: 238	SWISS: Q9UM73	Rabbit, Cow, Dog)
Target: ALK		
Immunogen: KLH conjugated synthetic peptide derived from human ALK: 19-120/1620. < Extracellular >		Predicted 174 kDa
Purification: affinity purified by	Protein A	
Concentration: 1mg/ml		Subcellular Location:
Storage: 0.01M TBS (pH7.4) Glycerol. Shipped at 4°C. St freeze/thaw cycles	with 1% BSA, 0.02% Proclin300 and 50% ore at -20°C for one year. Avoid repeated 5.	
Background: This gene encodes insulin receptor su extracellular dom single pass transm domain. It plays al and exerts its effec This gene has bee in a series of tumo neuroblastoma, al rearrangements a gene, which result tumourigenesis, ir (chromosome 2), / (chromosome 10), (chromosome 19), RefSeq, Jan 2011]	a receptor tyrosine kinase, which belong perfamily. This protein comprises an ain, an hydrophobic stretch corresponding nembrane region, and an intracellular kina n important role in the development of the cts on specific neurons in the nervous syst n found to be rearranged, mutated, or am urs including anaplastic large cell lympho nd non-small cell lung cancer. The chromo re the most common genetic alterations in including ALK (chromosome 2), ALK/ATIC ALK/TFG (chromosome 3), ALK/NPM1 ALK/SQSTM1 (chromosome 17), ALK/TPM4 , ALK/MSN (chromosome X).[provided	s to the g to a ase e brain em. .plified omas, osomal n this

- VALIDATION IMAGES



Sample: Cerebrum (Mouse) Lysate at 40 ug Primary: Anti-ALK (bs-23231R) at 1/300 dilution Secondary: IRDye800CW Goat Anti-Rabbit IgG at 1/20000 dilution Predicted band size: 174 kD Observed band size: 174 kD

- SELECTED CITATIONS -------

- [IF=5.168] Ying Gao. et al. ZYY-B-2, a novel ALK inhibitor, overcomes resistance to ceritinib by inhibiting P-gp function and induces apoptosis through mitochondrial pathway in ceritinib-resistant H2228 cells. CHEM-BIOL INTERACT. 2023 Jul;379:110516 WB ;Human. 37116853
- [IF=3.606] Xuejiao Zhou. et al. The novel ALK inhibitor ZX 29 induces apoptosis through inhibiting ALK and inducing ROS mediated endoplasmic reticulum stress in Karpas299 cells. 2020 Nov 02 WB ;Human. 33140567

• [IF=3.3] Yuying Yang. et al. EML4-ALK G1202R and EML4-ALK L1196M mutations induce crizotinib resistance in non-small cell lung cancer cells through activating epithelial-mesenchymal transition mediated by MDM2/MEK/ERK signal axis. CELL BIOL INT. 2024 Sep;: WB ;Human. 39318039