## bs-7648R

## [ Primary Antibody ]


## DFFB Rabbit pAb



www.bioss.com.cn sales@bioss.com.cn techsupport@bioss.com.cn 400-901-9800

| - DATASHEET  |               | 400-901-9800   |
|--|---------------|--|
| Host: Rabbit   | Isotype: IgG  | Applications: IHC-P (1:100-500)                                  |
| Clonality: Polyclonal  |               | IHC-F (1:100-500)<br>IF (1:100-500)                              |
| <b>GenelD:</b> 1677  | SWISS: 076075 |  |
| Target: DFFB   |               | <b>Reactivity:</b> Mouse (predicted: Human,<br>Rat, Rabbit, Dog) |
| Immunogen: KLH conjugated synthetic peptide derived from human DFFB: 251-338/338.  |               |  |
| Purification: affinity purified by Protein A   |               | Predicted<br>MW.: <sup>39 kDa</sup>                              |
| Concentration: 1mg/ml  |               |  |
| <b>Storage:</b> 0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50%<br>Glycerol.<br>Shipped at 4°C. Store at -20°C for one year. Avoid repeated<br>freeze/thaw cycles.   |               | Subcellular<br>Location: Cytoplasm ,Nucleus                      |
| <b>Background:</b> Apoptosis is a cell death process that removes toxic and/or useless cells during mammalian development. The apoptotic process is accompanied by shrinkage and fragmentation of the cells and nuclei and degradation of the chromosomal DNA into nucleosomal units. DNA fragmentation factor (DFF) is a heterodimeric protein of 40-kD (DFFB) and 45-kD (DFFA) subunits. DFFA is the substrate for caspase-3 and triggers DNA fragmentation during apoptosis. DFF becomes activated when DFFA is cleaved by caspase-3. The cleaved fragments of DFFA dissociate from DFFB, the active component of DFF. DFFB has been found to trigger both DNA fragmentation and chromatin condensation during apoptosis. Alternatively spliced transcript variants encoding distinct isoforms have been found for this gene but the biological validity of these variants has not been determined. [provided by RefSeq, Jul 2008]. |               |  |

## - VALIDATION IMAGES -



Paraformaldehyde-fixed, paraffin embedded (Mouse brain); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Antibody incubation with (DFFB) Polyclonal Antibody, Unconjugated (bs-7648R) at 1:400 overnight at 4°C, followed by operating according to SP Kit(Rabbit) (sp-0023) instructionsand DAB staining.