

bs-0452R**[Primary Antibody]****Bioss**
ANTIBODIES

www.bioss.com.cn


sales@bioss.com.cn

techsupport@bioss.com.cn

400-901-9800

NAIF1 Rabbit pAb**— DATASHEET —**

Host: Rabbit	Isotype: IgG	Applications: IHC-P (1:100-500) IHC-F (1:100-500) IF (1:100-500) Reactivity: Human (predicted: Mouse, Rat, Cow, Chicken, Dog, Horse) Predicted MW.: 35 kDa Subcellular Location: Nucleus
Clonality: Polyclonal		
GeneID: 203245	SWISS: Q69YI7	
Target: NAIF1		
Immunogen: KLH conjugated synthetic peptide derived from human NAIF1: 3-85/327.		
Purification: affinity purified by Protein A		
Concentration: 1mg/ml		
Storage: 0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol. Shipped at 4°C. Store at -20°C for one year. Avoid repeated freeze/thaw cycles.		
Background: NAIF protein 1 (nuclear apoptosis inducing factor 1; CB12-327; RP-379C10.2) belongs to the NAIF1 family. Apoptosis is a genetically determined cell suicidal program that plays critical roles in many physiological and pathological processes. nuclear apoptosis-inducing factor 1 (NAIF1), overexpression of which induces apoptosis in cells. Human NAIF1 is located on chromosome 9q34.11 and encodes 327 amino acids with a homeodomain-like region and two nuclear localization signals at its N-terminal region. NAIF1 is conserved across diverse species, including human, mouse, crab-eating macaque, dog, chicken and frog, and shares no obvious homology to any known genes or proteins. Northern blot analysis revealed wide expression of NAIF1 mRNA throughout human tissues. NAIF1 was predominantly localized in the nucleus. Overexpression of NAIF1 inhibited cell growth and induced apoptosis. Furthermore, NAIF1 transfection caused both decreases in mitochondrial membrane potential and caspase-3 activation. In summary, NAIF1 is a nuclear protein that induces apoptosis when overexpressed.		

— VALIDATION IMAGES —

Tissue/cell: human colon tissue; 4% Paraformaldehyde-fixed and paraffin-embedded; Antigen retrieval: citrate buffer (0.01M, pH 6.0), Boiling bathing for 15min; Block endogenous peroxidase by 3% Hydrogen peroxide for 30min; Blocking buffer (normal goat serum, C-0005) at 37°C for 20 min; Incubation: Anti-NAIF1 Polyclonal Antibody, Unconjugated(bs-0452R) 1:200, overnight at 4°C, followed by conjugation to the secondary antibody(SP-0023) and DAB(C-0010) staining