bs-5389R

[Primary Antibody]

phospho-HDAC2 (Ser394) Rabbit pAb



www.bioss.com.cn sales@bioss.com.cn techsupport@bioss.com.cn 400-901-9800

DATASHEET -

Host: Rabbit Isotype: IgG

Clonality: Polyclonal

GeneID: 3066 SWISS: Q92769

Target: HDAC2 (Ser394)

Immunogen: KLH conjugated Synthesised phosphopeptide derived from human

HDAC2 around the phosphorylation site of Ser394: ED(p-S)GD.

Purification: affinity purified by Protein A

Concentration: 1mg/ml

Storage: 0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50%

Glycerol.

Shipped at 4°C. Store at -20°C for one year. Avoid repeated

freeze/thaw cycles.

Background: This gene product belongs to the histone deacetylase family. Histone deacetylases act via the formation of large multiprotein complexes, and are responsible for the deacetylation of lysine residues at the N-terminal regions of core histones (H2A, H2B, H3 and H4). This protein forms transcriptional repressor complexes by associating with many different proteins, including YY1, a mammalian zinc-finger transcription factor. Thus, it plays an important role in transcriptional regulation, cell cycle progression and developmental events. Alternative splicing results in multiple transcript variants. [provided by RefSeq].

Applications: WB (1:500-2000)

IHC-P (1:100-500) IHC-F (1:100-500) **IF** (1:100-500) Flow-Cyt (2ug/Test) ICC/IF (1:100-500)


Reactivity: Human, Mouse, Rat

(predicted: Pig, Cow, Chicken, Dog, Horse)


Predicted MW.: 55 kDa

Subcellular Location: Nucleus


VALIDATION IMAGES




Sample: Lane 1: Mouse Cerebrum tissue lysates Lane 2: Mouse NIH/3T3 cell lysates Lane 3: Rat Cerebrum tissue lysates Primary: Anti-phospho-HDAC2 (Ser394) (bs-5389R) at 1/1000 dilution Secondary: IRDve800CW Goat Anti-Rabbit IgG at 1/20000 dilution Predicted band size: 55 kDa Observed band size: 52 kDa



Paraformaldehyde-fixed, paraffin embedded (mouse intestine tissue); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes: Blocking buffer (normal goat serum) at 37°C for 30min; Antibody incubation with (phospho-HDAC2 (Ser394)) Polyclonal Antibody, Unconjugated (bs-5389R) at 1:400 overnight at 4°C, followed by operating according to SP Kit(Rabbit) (sp-0023) instructions and DAB staining.



Hela cell; 4% Paraformaldehyde-fixed; Triton X-100 at room temperature for 20 min; Blocking buffer (normal goat serum, C-0005) at 37°C for 20 min; Antibody incubation with (phospho-HDAC2 (Ser394)) polyclonal Antibody, Unconjugated (bs-5389R) 1:50, 90 minutes at 37°C; followed by a conjugated Goat Anti-Rabbit IgG antibody at 37°C for 90 minutes, DAPI (blue, C02-04002) was used to stain the cell nuclei.



Blank control (black line) :Hela, Primary Antibody (green line): Rabbit Anti-phospho-HDAC2(Ser394) antibody (bs-5389R) Dilution:2ug/Test; Secondary Antibody (white blue line): Goat anti-rabbit IgG-FITC Dilution: 0.5ug/Test. Isotype control (orange line): Normal Rabbit IgG Protocol The cells were fixed with 4% PFA (10min at room temperature) and then permeabilized with 90% ice-cold methanol for 20 min at -20°C, The cells were then incubated in 5%BSA to block non-specific protein-protein interactions for 30 min at room temperature. Cells stained with Primary Antibody for 30 min at room temperature. The secondary antibody used for 40 min at room temperature. Acquisition of 20,000 events was performed.

- SELECTED CITATIONS -

• [IF=3.81] Jou, Yu - Jen, et al. "Quantitative phosphoproteomic analysis reveals γ-bisabolene inducing p53 - mediated apoptosis of human oral squamous cell carcinoma via HDAC2 inhibition and ERK1/2 activation." Proteomics (2015). WB; Human. 26194454