
TRAK1 Rabbit pAb

Catalog Number: bs-5536R

Target Protein: TRAK1

Concentration: 1mg/ml

Form: Liquid

Host: Rabbit

Clonality: Polyclonal

Isotype: IgG

Applications: **IHC-P** (1:100-500), **IHC-F** (1:100-500), **IF** (1:100-500), **ICC/IF** (1:100)

Reactivity: Human, Mouse (predicted: Rat, Pig, Cow, Chicken, Dog, Horse)

Predicted MW: 106 kDa

Subcellular: Cytoplasm, Nucleus

Locations:

Entrez Gene: 22906

Swiss Prot: Q9UPV9

Source: KLH conjugated synthetic peptide derived from human TRAK1: 171-270/953.


Purification: affinity purified by Protein A

Storage: 0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol.


Shipped at 4°C. Store at -20°C for one year. Avoid repeated freeze/thaw cycles.

Background: The specific function of TRAK1 remains unknown. It interacts with O-GlcNAc transferase, RHOT1/Miro-1 and RHOT2/Miro-2. It shows high expression in spinal cord and moderate expression in all other tissues and specific brain regions examined.


VALIDATION IMAGES




Paraformaldehyde-fixed, paraffin embedded (human gastric carcinoma); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Antibody incubation with (TRAK1) Polyclonal Antibody, Unconjugated (bs-5536R) at 1:200 overnight at 4°C, followed by operating according to SP Kit(Rabbit) (sp-0023) instructions and DAB staining.



Paraformaldehyde-fixed, paraffin embedded (human liver cancer); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Antibody incubation with (TRAK1) Polyclonal Antibody, Unconjugated (bs-5536R) at 1:200 overnight at 4°C, followed by operating according to SP Kit(Rabbit) (sp-0023) instructions and DAB staining.



Paraformaldehyde-fixed, paraffin embedded (mouse brain tissue); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Antibody incubation with (TRAK1) Polyclonal Antibody, Unconjugated (bs-5536R) at 1:200 overnight at 4°C, followed by operating according to SP Kit(Rabbit) (sp-0023) instructions and DAB staining.



Hela cell; 4% Paraformaldehyde-fixed; Triton X-100 at room temperature for 20 min; Blocking buffer (normal goat serum, C-0005) at 37°C for 20 min; Antibody incubation with (TRAK1) polyclonal Antibody, Unconjugated (bs-5536R) 1:100, 90 minutes at 37°C; followed by a conjugated Goat Anti-Rabbit IgG antibody at 37°C for 90 minutes, DAPI (blue, C02-04002) was used to stain the cell nuclei.