

bs-2139R**[Primary Antibody]****GRM2+GRM4 Rabbit pAb****BioSS**
ANTIBODIES

www.bioss.com.cn


sales@bioss.com.cn

techsupport@bioss.com.cn

400-901-9800

— DATASHEET —

<p>Host: Rabbit</p> <p>Clonality: Polyclonal</p> <p>GeneID: 2912</p> <p>Target: GRM2+GRM4</p> <p>Immunogen: KLH conjugated synthetic peptide derived from human GRM2+GRM4: 31-130/872.</p> <p>Purification: affinity purified by Protein A</p> <p>Concentration: 1mg/ml</p> <p>Storage: 0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol. Shipped at 4°C. Store at -20°C for one year. Avoid repeated freeze/thaw cycles.</p> <p>Background: L-glutamate is the major excitatory neurotransmitter in the central nervous system and activates both ionotropic and metabotropic glutamate receptors. Glutamatergic neurotransmission is involved in most aspects of normal brain function and can be perturbed in many neuropathologic conditions. The metabotropic glutamate receptors are a family of G protein-coupled receptors, that have been divided into 3 groups on the basis of sequence homology, putative signal transduction mechanisms, and pharmacologic properties. Group I includes GRM1 and GRM5 and these receptors have been shown to activate phospholipase C. Group II includes GRM2 and GRM3 while Group III includes GRM4, GRM6, GRM7 and GRM8. Group II and III receptors are linked to the inhibition of the cyclic AMP cascade but differ in their agonist selectivities.</p>	<p>Isotype: IgG</p> <p>SWISS: Q14416</p> <p>Applications: WB (1:500-2000)</p> <p>Reactivity: Human (predicted: Mouse, Rat)</p> <p>Predicted MW.: 102 kDa</p> <p>Subcellular Location: Cell membrane</p>
--	---

— VALIDATION IMAGES —

Sample: Lane 1: Human HepG2 cell lysates Lane 2: Human HeLa cell lysates Lane 3: Human K562 cell lysates Lane 4: Human SH-SY5Y cell lysates
 Primary: Anti-GRM2+GRM4 (bs-2139R) at 1/1000 dilution Secondary: IRDye800CW Goat Anti-Rabbit IgG at 1/20000 dilution Predicted band size: 102 kDa Observed band size: 120 kDa