bs-5398R

[Primary Antibody]

BIOSS ANTIBODIES

www.bioss.com.cn sales@bioss.com.cn techsupport@bioss.com.cn 400-901-9800

phospho-IKK beta (Ser471) Rabbit pAb

- DATASHEET -

Host: Rabbit Isotype: IgG

Clonality: Polyclonal

GenelD: 3551 **SWISS:** 014920

Target: IKK beta (Ser471)

Immunogen: KLH conjugated Synthesised phosphopeptide derived from human

IKK beta around the phosphorylation site of Ser471: KN(p-S)MA.

Purification: affinity purified by Protein A

Concentration: 1mg/ml

Storage: 0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50%

Glycerol.

Shipped at 4°C. Store at -20°C for one year. Avoid repeated

freeze/thaw cycles.

Background: IKK beta (I-Kappa-B kinase-beta) is a member of the IKK complex

which is composed of IKK alpha, IKK beta, IKK gamma and IKAP. Phosphorylation of I-Kappa-B on a serine residue by the IKK complex frees NF-kB from I-Kappa-B and marks it for degradation via ubiquination. IKK beta has been shown to activate NF-kB and phosphorylate IKB alpha and beta. Phosphorylation of 2 sites at the activation loop of IKK beta is essential for activation of IKK by TNF and IL1. Once activated, IKK beta autophosphorylates which in turn decreases IKK activity and prevents prolonged activation of the inflammatory response. Additionally, IKK beta activity can also

be regulated by MEKK1.

Applications: IHC-P (1:100-500)

IHC-F (1:100-500) IF (1:100-500)

Reactivity: Mouse (predicted: Human,


Rat, Rabbit, Pig, Cow, Dog)

Predicted MW • 87 kD

Subcellular Cell membrane ,Cytoplasm

Location: , Nucleus

VALIDATION IMAGES



Paraformaldehyde-fixed, paraffin embedded (Mouse placenta); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Antibody incubation with (phospho-IKK beta(Ser471)) Polyclonal Antibody, Unconjugated (bs-5398R) at 1:400 overnight at 4°C, followed by operating according to SP Kit(Rabbit) (sp-0023) instructionsand DAB staining.

SFI FCTFD CITATIONS -

• [IF=2.447] Mingdong Si. et al. Gentianella acuta mitigates cardiovascular damage and inflammation in diet-induced hypercholesterolaemic rats. Exp Ther Med. 2021 Nov;22(5):1-10 WB,IHC;rat. 34603527