bs-5399R

## [ Primary Antibody ]

## phospho-IKK beta (Ser740) Rabbit pAb



www.bioss.com.cn sales@bioss.com.cn techsupport@bioss.com.cn 400-901-9800

DATASHEET -

Host: Rabbit Isotype: IgG

Clonality: Polyclonal

**GenelD:** 3551 **SWISS:** 014920

Target: IKK beta (Ser740)

Immunogen: KLH conjugated Synthesised phosphopeptide derived from human

IKK beta around the phosphorylation site of Ser740: DW(p-S)WL.

**Purification:** affinity purified by Protein A

Concentration: 1mg/ml

**Storage:** 0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50%

Glycerol.

Shipped at 4°C. Store at -20°C for one year. Avoid repeated

freeze/thaw cycles.

**Background:** IKK beta (I-Kappa-B kinase-beta) is a member of the IKK complex

which is composed of IKK alpha, IKK beta, IKK gamma and IKAP. Phosphorylation of I-Kappa-B on a serine residue by the IKK complex frees NF-kB from I-Kappa-B and marks it for degradation via ubiquination. IKK beta has been shown to activate NF-kB and phosphorylate IKB alpha and beta. Phosphorylation of 2 sites at the activation loop of IKK beta is essential for activation of IKK by TNF and IL1. Once activated, IKK beta autophosphorylates which in turn decreases IKK activity and prevents prolonged activation of the inflammatory response. Additionally, IKK beta activity can also

be regulated by MEKK1.

Applications: ICC/IF (1:50-200)

Reactivity: Human (predicted: Mouse,


Rat, Chicken)

Predicted MW.: 87 kDa

Subcellular Cell membrane ,Cytoplasm

Location: , Nucleus

## VALIDATION IMAGES



A549 cell; 4% Paraformaldehyde-fixed; Triton X-100 at room temperature for 20 min; Blocking buffer (normal goat serum, C-0005) at 37°C for 20 min; Antibody incubation with (phospho-IKK beta (Ser740)) polyclonal Antibody, Unconjugated (bs-5399R) 1:100, 90 minutes at 37°C; followed by a conjugated Goat Anti-Rabbit IgG antibody at 37°C for 90 minutes, DAPI (blue, C02-04002) was used to stain the cell nuclei.

## - SELECTED CITATIONS -

- [IF=6.117] Yuanyuan Xing. et al. Artemisia ordosica polysaccharide ameliorated LPS-induced growth inhibition and intestinal injury in broilers through enhancing immune-regulation and antioxidant capacity. J NUTR BIOCHEM. 2023 Feb;:109284 WB ;Chicken. 36828238
- [IF=6.1] Dongxue Song. et al. Purple Sweet Potato Polysaccharide Exerting an Anti-inflammatory Effect via a TLR-Mediated Pathway by Regulating Polarization and Inhibiting the Inflammasome Activation. J AGR FOOD CHEM. 2024;XXXX(XXX):XXX-XXX WB; Mouse. 38233194
- [IF=3.1] Zhou Shunhua. et al. Atractylenolide III as a novel therapeutic strategy for gastric ulcer: mechanistic insights

and investigation o	f molecular targets. N-:	S ARCH PHARMACU	L. 2025 Jul;:1-18 IF	,Kat. 40728671	