

bs-2184R**[Primary Antibody]****NQO1 Rabbit pAb****Bioss**
ANTIBODIES

www.bioss.com.cn


sales@bioss.com.cn

techsupport@bioss.com.cn


400-901-9800

— DATASHEET —**Host:** Rabbit**Isotype:** IgG**Clonality:** Polyclonal**GeneID:** 1728**SWISS:** P15559**Target:** NQO1**Immunogen:** KLH conjugated synthetic peptide derived from human NQO1: 201-274/274.**Purification:** affinity purified by Protein A**Concentration:** 1mg/ml**Storage:** 0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol.


Shipped at 4°C. Store at -20°C for one year. Avoid repeated freeze/thaw cycles.

Background: This gene is a member of the NAD(P)H dehydrogenase (quinone) family and encodes a cytoplasmic 2-electron reductase. This FAD-binding protein forms homodimers and reduces quinones to hydroquinones. This protein's enzymatic activity prevents the one electron reduction of quinones that results in the production of radical species. Mutations in this gene have been associated with tardive dyskinesia (TD), an increased risk of hematotoxicity after exposure to benzene, and susceptibility to various forms of cancer. Altered expression of this protein has been seen in many tumors and is also associated with Alzheimer's disease (AD). Alternate transcriptional splice variants, encoding different isoforms, have been characterized. [provided by RefSeq].**Applications:** **WB** (1:500-2000)**IHC-P** (1:100-500)**IHC-F** (1:100-500)**IF** (1:100-500)**ICC/IF** (1:50-200)**Reactivity:** Human, Mouse, Rat
(predicted: Rabbit, Pig, GuineaPig, Horse)**Predicted MW.:** 31 kDa**Subcellular Location:** Cytoplasm**— VALIDATION IMAGES —**


Sample: A549(Human) Cell Lysate at 30 ug
 Primary: Anti-NQO1 (bs-2184R) at 1/1000
 dilution Secondary: IRDye800CW Goat Anti-Rabbit IgG at 1/20000 dilution Predicted band size: 31 kD Observed band size: 31 kD




25 ug total protein per lane of various lysates (see on figure) probed with NQO1 polyclonal antibody, unconjugated (bs-2184R) at 1:1000 dilution and 4°C overnight incubation. Followed by conjugated secondary antibody incubation at r.t. for 60 min.



Tissue/cell: mouse colon carcinoma; 4% Paraformaldehyde-fixed and paraffin-embedded; Antigen retrieval: citrate buffer (0.01M, pH 6.0), Boiling bathing for 15min; Block endogenous peroxidase by 3% Hydrogen peroxide for 30min; Blocking buffer (normal goat serum, C-0005) at 37°C for 20 min; Incubation: Anti-NQO1 Polyclonal Antibody, Unconjugated(bs-2184R) 1:200, overnight at 4°C, followed by conjugation to the secondary antibody(SP-0023) and DAB(C-0010) staining



Paraformaldehyde-fixed, paraffin embedded (rat stomach); Antigen retrieval by boiling in sodium



4% Paraformaldehyde-fixed HepG2 (H) cell; Triton X-100 at r.t. for 20 min; Antibody

Important Note: This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.

citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Antibody incubation with (NQO1) Polyclonal Antibody, Unconjugated (bs-2184R) at 1:200 overnight at 4°C, followed by operating according to SP Kit(Rabbit) (sp-0023) instructions and DAB staining.

incubation with (NQO1) polyclonal Antibody, unconjugated (bs-2184R) 1:100, 90 min at 37°C; followed by conjugated Goat Anti-Rabbit IgG antibody (green, bs-40295G-FITC) at 37°C for 90 min, DAPI (blue, C02-04002) was used to stain the cell nuclei. PBS instead of the primary antibody was used as the blank control.

— SELECTED CITATIONS —

- **[IF=15.1]** Wenting Li. et al. A collective hyperthermia-facilitated glutathione inhibition strategy for boosted oxidative stress treatment. CHEM ENG J. 2024 Apr;485:150056 WB ;Mouse. 10.1016/j.cej.2024.150056
- **[IF=7.9]** Yu-Sheng Shi. et al. Dendrobine rescues cognitive dysfunction in diabetic encephalopathy by inhibiting ferroptosis via activating Nrf2/GPX4 axis. PHYTOMEDICINE. 2023 Jul;154993 WB ;Mouse. 37567006
- **[IF=8.2]** Yaqi Li. et al. Non-thermal plasma promotes boar sperm quality through increasing AMPK methylation. INT J BIOL MACROMOL. 2024 Feb;257:128768 WB ;Pig. 38096931
- **[IF=7.129]** Rui Sang. et al. Taraxasterol alleviates aflatoxin B1-induced liver damage in broiler chickens via regulation of oxidative stress, apoptosis and autophagy. ECOTOX ENVIRON SAFE. 2023 Feb;251:114546 WB ;Chicken. 36646010
- **[IF=6.953]** Jinshan Wu. et al. Protection by Hosta ventricosa polysaccharides against oxidative damage induced by t-BHP in HepG2 cells via the JNK/Nrf2 pathway. Int J Biol Macromol. 2022 May;208:453 WB ;Human. 35339497