

bs-6474R**[Primary Antibody]****connexin 30 Rabbit pAb****BioSS**
ANTIBODIES

www.bioss.com.cn


sales@bioss.com.cn

techsupport@bioss.com.cn

400-901-9800

— DATASHEET —

Host: Rabbit	Isotype: IgG	Applications: IHC-P (1:100-500) IHC-F (1:100-500) IF (1:50-200)
Clonality: Polyclonal		
GeneID: 10804	SWISS: Q95452	
Target: connexin 30		Reactivity: Rat (predicted: Human, Mouse, Sheep, Cow, Chicken, Dog, Horse)
Immunogen: KLH conjugated synthetic peptide derived from human connexin-30: 161-261/260. < Extracellular >		
Purification: affinity purified by Protein A		Predicted MW.: 29 kDa
Concentration: 1mg/ml		Subcellular Location: Cell membrane
Storage: 0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol. Shipped at 4°C. Store at -20°C for one year. Avoid repeated freeze/thaw cycles.		
Background: The connexin family of proteins form hexameric complexes called connexons that facilitate movement of low molecular weight proteins between cells via gap junctions. Connexin proteins share a common topology of four transmembrane alpha-helical domains, two extracellular loops, a cytoplasmic loop and cytoplasmic N- and C-termini. Many of the key functional differences between connexins arise from specific amino-acid substitutions in the most highly conserved domains: the transmembrane and extracellular regions. Connexin 30, also known as GJB6 (Gap junction beta 6), ED2, EDH, HED or DFNA3, is a 261 amino acid multi-pass membrane protein that localizes to the cell junction and belongs to the connexin family. Functioning as a hexamer with other connexin proteins, connexin 30 facilitates the diffusion of low molecular weight materials from one cell to another. Defects in the gene encoding connexin 30 are the cause of ectodermal dysplasia type 2 (ED2) and non-syndromic sensorineural deafness autosomal dominant type 3 (DFNA3), the former of which is characterized by abnormal development of ectodermal structures (such as skin and nails).		

— VALIDATION IMAGES —

Tissue/cell: Rat tongue; 4% Paraformaldehyde-fixed and paraffin-embedded; Antigen retrieval: citrate buffer (0.01M, pH 6.0), Boiling bathing for 15min; Block endogenous peroxidase by 3% Hydrogen peroxide for 30min; Blocking buffer (normal goat serum,C-0005) at 37°C for 20 min; Incubation: Anti-connexin 30/GJB6 Polyclonal Antibody, Unconjugated (bs-6474R) 1:200, overnight at 4°C, followed by conjugation to the secondary antibody(SP-0023) and DAB(C-0010) staining