

www.bioss.com.cn sales@bioss.com.cn techsupport@bioss.com.cn 400-901-9800

## MARCH9 Rabbit pAb

Catalog Number: bs-9343R
Target Protein: MARCH9

Concentration: 1mg/ml

Form: Liquid

Host: Rabbit

Clonality: Polyclonal

Isotype: IgG

Applications: WB (1:500-2000), Flow-Cyt (1µg/Test)

Reactivity: Human (predicted: Mouse, Rat, Rabbit, Pig, Sheep, Cow, Chicken, Dog, Horse)

Predicted MW: 38 kDa Entrez Gene: 92979 Swiss Prot: Q86YJ5

**Source:** KLH conjugated synthetic peptide derived from human MARCH9: 201-300/346.

Purification: affinity purified by Protein A


Storage: 0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol.

Shipped at 4°C. Store at -20°C for one year. Avoid repeated freeze/thaw cycles.


Background: Ubiquitination is an important mechanism through which three classes of enzymes act in

concert to target short-lived or abnormal proteins for destruction. The three classes of enzymes involved in ubiquitination are the ubiquitin-activating enzymes (E1s), the ubiquitin-conjugating enzymes (E2s) and the ubiquitin-protein ligases (E3s). MARCH9 (membrane-associated ring finger (C3HC4) 9), also known as RNF179, is a 346 amino acid multi-pass membrane protein that localizes to the golgi apparatus and contains one RING-CH-type zinc finger. Expressed ubiquitously, MARCH9 exists as a homodimer and functions as an E3 ubiquitin-protein ligase that accepts a ubiquitin residue from an E2 ubiquitin-conjugating enzyme and is thought to promote the degradation of target proteins, such as CD4 and MHC-I. Multiple isoforms of MARCH9 exist due to alternative splicing events.

**VALIDATION IMAGES** 



Sample: MCF-7 (human)cell Lysate at 40 ug Primary: Anti- MARCH9(bs-9343R)at 1/300 dilution Secondary: IRDye800CW Goat Anti-Rabbit IgG at 1/20000 dilution Predicted band size: 38kD Observed band size: 38 kD



Blank control: Hela(blue), the cells were fixed with 2% paraformaldehyde (10 min) Isotype Control Antibody: Rabbit IgG(orange); Secondary Antibody: Goat anti-rabbit IgG-FITC(white blue), Dilution: 1:200 in 1 X PBS containing 0.5% BSA; Primary Antibody Dilution:  $1\mu g$  in 100  $\mu L$  1X PBS containing 0.5% BSA(green).