

bs-5711R**[Primary Antibody]****CCDC106 Rabbit pAb****Bioss**
ANTIBODIES

www.bioss.com.cn


sales@bioss.com.cn

techsupport@bioss.com.cn


400-901-9800

— DATASHEET —


Host: Rabbit	Isotype: IgG	Applications: WB (1:500-2000) IHC-P (1:100-500) IHC-F (1:100-500) IF (1:100-500)
Clonality: Polyclonal		
GeneID: 29903	SWISS: Q9BWC9	
Target: CCDC106		
Immunogen: KLH conjugated synthetic peptide derived from human CCDC106: 201-280/280.		
Purification: affinity purified by Protein A		
Concentration: 1mg/ml		Reactivity: Human, Mouse, Rat (predicted: Cow, Dog, Horse)
Storage: 0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol. Shipped at 4°C. Store at -20°C for one year. Avoid repeated freeze/thaw cycles.		Predicted MW.: 31 kDa
Background: Promotes the degradation of p53/TP53 protein and inhibits its transactivity.		Subcellular Location: Nucleus

— VALIDATION IMAGES —


Sample: Jurkat(Human) Cell Lysate at 30 ug
 Primary: Anti- CCDC106 (bs-5711R) at 1/1000 dilution
 Secondary: IRDye800CW Goat Anti-Rabbit IgG at 1/20000 dilution
 Predicted band size: 31 kD Observed band size: 31 kD



Sample: A549(Human) Cell Lysate at 30 ug
 Primary: Anti- CCDC106 (bs-5711R) at 1/1000 dilution
 Secondary: IRDye800CW Goat Anti-Rabbit IgG at 1/20000 dilution
 Predicted band size: 31 kD Observed band size: 31 kD



Paraformaldehyde-fixed, paraffin embedded (mouse brain); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Antibody incubation with (CCDC106) Polyclonal Antibody, Unconjugated (bs-5711R) at 1:200 overnight at 4°C, followed by operating according to SP Kit(Rabbit) (sp-0023) instructions and DAB staining.



Paraformaldehyde-fixed, paraffin embedded (rat brain); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Antibody incubation with (CCDC106) Polyclonal Antibody, Unconjugated (bs-5711R) at 1:200 overnight at 4°C, followed by operating according to SP Kit(Rabbit) (sp-0023) instructions and DAB staining.

Important Note: This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.

4°C, followed by operating according to SP
Kit(Rabbit) (sp-0023) instructions and DAB
staining.