## [ Primary Antibody ]

## Scavenger Receptor BII Rabbit pAb



www.bioss.com.cn sales@bioss.com.cn techsupport@bioss.com.cn 400-901-9800

		400-901-9800
Host: Rabbit	Isotype: IgG	Applications: WB (1:500-2000)
Clonality: Polyclonal		<b>Poactivity:</b> Human (prodicted: Mouse
GenelD: 950	SWISS: Q14108	Rat, Rabbit, Pig, Cow, Dog,
Target: Scavenger Receptor	BII	Horse)
Immunogen: KLH conjugated synthetic peptide derived from human Scavenger Receptor BII: 181-280/478.		Predicted 52 kDa
Purification: affinity purified by Protein A		
Concentration: 1mg/ml		Subcellular Location: Cell membrane ,Cytoplasm
<b>Storage:</b> 0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol. Shipped at 4°C. Store at -20°C for one year. Avoid repeated freeze/thaw cycles.		
<b>Background:</b> High density lipoproteins (HDLs) play a critical role in cholesterol metabolism and their plasma concentrations are inversely correlated with risk for atherosclerosis. SR-BI and SR-BII (previously known as SR-BI.2) are the alternatively spliced products of a single gene. SR-BI and SR-BI are identical except for the encoded c-terminal cytoplasmic domain. Both SR-BI and SR-BII bind HDL and mediates selective uptake of HDL cholesteryl ester, but with SR-BII having an approximately 4-fold lower efficiency than SR-BI. SR-BI and SR-BII are expressed primarily in liver and non-placental steroidgenic tissues. Although the role of these scavenger receptors is not completely clear, SR-BII mRNA results from the alternative splicing of SR-BI precursor transcripts with both isoforms mediating selective transfer of lipid between HDL and cells. Therefore, the relative expression and functional activities of these two isoforms create a potential means of		

## - VALIDATION IMAGES -----



Sample: Hela Cell (Human) Lysate at 40 ug Primary: Anti-Scavenger Receptor BII (bs-7545R) at 1/300 dilution Secondary: IRDye800CW Goat Anti-Rabbit IgG at 1/20000 dilution Predicted band size: 52 kD Observed band size: 52 kD