

bs-3097R**[Primary Antibody]****phospho-CDK9 (Thr186) Rabbit pAb****Bioss**
ANTIBODIES

www.bioss.com.cn


sales@bioss.com.cn

techsupport@bioss.com.cn


400-901-9800

DATASHEET**Host:** Rabbit**Isotype:** IgG**Clonality:** Polyclonal**GeneID:** 1025**SWISS:** P50750**Target:** CDK9 (Thr186)**Immunogen:** KLH conjugated Synthesised phosphopeptide derived from human CDK9 around the phosphorylation site of Thr186: RY(p-T)NR.**Purification:** affinity purified by Protein A**Concentration:** 1mg/ml**Storage:** 0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol.


Shipped at 4°C. Store at -20°C for one year. Avoid repeated freeze/thaw cycles.

Background: The protein encoded by this gene belongs to the highly conserved cyclin family, whose members are characterized by a dramatic periodicity in protein abundance through the cell cycle. Cyclins function as regulators of CDK kinases. Different cyclins exhibit distinct expression and degradation patterns which contribute to the temporal coordination of each mitotic event. This cyclin tightly associates with CDK9 kinase, and was found to be a major subunit of the transcription elongation factor p-TEFb. The kinase complex containing this cyclin and the elongation factor can interact with, and act as a cofactor of human immunodeficiency virus type 1 (HIV-1) Tat protein, and was shown to be both necessary and sufficient for full activation of viral transcription. This cyclin and its kinase partner were also found to be involved in the phosphorylation and regulation of the carboxy-terminal domain (CTD) of the largest RNA polymerase II subunit.**Applications:** WB (1:500-2000)**IHC-P** (1:100-500)**IHC-F** (1:100-500)**IF** (1:100-500)**Reactivity:** Human, Mouse
(predicted: Rat, Pig, Cow, Chicken, Dog, Horse)**Predicted MW.:** 43 kDa**Subcellular Location:** Cytoplasm ,Nucleus**VALIDATION IMAGES**

Sample: Brain (Mouse) Lysate at 40 ug Primary: Anti-p-CDK9 (Thr186) (bs-3097R) at 1/300 dilution Secondary: IRDye800CW Goat Anti-Rabbit IgG at 1/20000 dilution Predicted band size: 43 kD Observed band size: 48 kD



Sample: Lane 1: Human 293T cell lysates Lane 2: Human MCF-7 cell lysates Lane 3: Human HepG2 cell lysates Lane 4: Human HeLa cell lysates Lane 5: Human Jurkat cell lysates Lane 6: Human A431 cell lysates Primary: Anti-Phospho-CDK9 (Thr186) (bs-3097R) at 1/500 dilution Secondary: IRDye800CW Goat Anti-Rabbit IgG at 1/20000 dilution Predicted band size: 43 kDa Observed band size: 42,55 kDa



Tissue/cell: human laryngeal tissue; 4% Paraformaldehyde-fixed and paraffin-embedded; Antigen retrieval: citrate buffer (0.01M, pH 6.0), Boiling bathing for 15min; Block endogenous peroxidase by 3% Hydrogen peroxide for 30min; Blocking buffer (normal goat serum, C-0005) at 37°C for 20 min; Incubation: Anti-Phospho-CDK9 (Thr186) Polyclonal Antibody, Unconjugated(bs-3097R) 1:200, overnight at 4°C, followed by conjugation to the secondary antibody(SP-0023) and DAB(C-0010) staining