
NPFF1 Receptor Rabbit pAb

Catalog Number: bs-12018R

Target Protein: NPFF1 Receptor

Concentration: 1mg/ml

Form: Liquid

Host: Rabbit

Clonality: Polyclonal

Isotype: IgG

Applications: WB (1:500-2000), IHC-P (1:100-500), IHC-F (1:100-500), IF (1:100-500)

Reactivity: Human, Mouse, Rat (predicted:Rabbit, Pig, Sheep, Cow, Dog, Horse)

Predicted MW: 48 kDa

Entrez Gene: 64106

Swiss Prot: Q9GZQ6

Source: KLH conjugated synthetic peptide derived from human GPR147/NPFF1 Receptor:
151-260/430.


Purification: affinity purified by Protein A

Storage: 0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol.


Shipped at 4°C. Store at -20°C for one year. Avoid repeated freeze/thaw cycles.

Background: Neuropeptide FF 1 Receptor (NPFF1 or hFF1) and Neuropeptide FF 2 Receptor (NPFF2) belong to the G protein-coupled receptor 1 family. Both NPFF1 and NPFF2 are integral membrane proteins that act as receptors for NPAF (A-18- F-amide) and NPFF (F-8-F-amide) neuropeptides. Both NPFF proteins may be activated by synthetic or naturally occurring FMRF-amide-like ligands. The receptors are mediated by association with G proteins that activate a phosphatidylinositol-calcium second messenger system. NPFF1 Receptors is highly expressed in the human hypothalamus and amygdala, indicating a possible role for NPFF1 in central autonomic and neuroendocrine control in the human brain. Based in part on NPFF2 Receptor expression in diencephalon and superficial layers of the spinal cord, NPFF2 Receptor is thought to be involved in the modulation of sensory input and opioid analgesia.


VALIDATION IMAGES




Sample: U251(Human) Cell Lysate at 30 ug Primary: Anti-NPFF1 Receptor (bs-12018R) at 1/300 dilution
Secondary: IRDye800CW Goat Anti-Rabbit IgG at 1/20000 dilution Predicted band size: 48 kD Observed band size: 48 kD



Sample: Lane 1: Mouse Cerebellum tissue lysates Lane 2: Mouse Cerebrum tissue lysates Lane 3: Rat Cerebellum tissue lysates Lane 4: Rat Cerebrum tissue lysates Primary: Anti-NPFF1 Receptor (bs-12018R) at 1/1000 dilution Secondary: IRDye800CW Goat Anti-Rabbit IgG at 1/20000 dilution Predicted band size: 48 kDa Observed band size: 48 kDa



Paraformaldehyde-fixed, paraffin embedded (Rat brain); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Antibody incubation with (NPFF1 Receptor) Polyclonal Antibody, Unconjugated (bs-12018R) at 1:400 overnight at 4°C, followed by operating according to SP Kit(Rabbit) (sp-0023) instructions and DAB staining.



Paraformaldehyde-fixed, paraffin embedded (Mouse brain); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Antibody incubation with (NPFF1 Receptor) Polyclonal Antibody, Unconjugated (bs-12018R) at 1:400 overnight at 4°C, followed by operating according to SP Kit(Rabbit) (sp-0023) instructions and DAB staining.

PRODUCT SPECIFIC PUBLICATIONS

[IF=3.333] Zixuan Chen. et al. Exogenous Melatonin Regulates Puberty and the Hypothalamic GnRH-GnIH System in Female Mice. BRAIN SCI. 2022 Nov;12(11):1550 WB ; Mouse . 36421874

[IF=2.751] Xueying Zhao. et al. Effects of RFRP-3 on an ovariectomized estrogen-primed rat model and HEC-1A human endometrial carcinoma cells. EXP THER MED. 2023 Feb;25(2):1-14 WB ; Human . 36684658