bs-17700R

[Primary Antibody]

MOCS1 Rabbit pAb



www.bioss.com.cn sales@bioss.com.cn techsupport@bioss.com.cn 400-901-9800

- DATASHEET		400-901-9800
Host: Rabbit	Isotype: IgG	Applications: IHC-P (1:100-500)
Clonality: Polyclonal		IHC-F (1:100-500)
GenelD: 4337	SWISS: Q9NZB8	ICC/IF (1:100-500)
Target: MOCS1		ELISA (1:5000-10000)
Immunogen: KLH conjugated synthetic peptide derived from human MOCS1: 21-120/636.		Reactivity: (predicted: Human, Mouse, Rat, Rabbit, Pig, Horse)
Purification: affinity purified by I	Protein A	
Concentration: 1mg/ml		Due diete d
Storage: 0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol. Shipped at 4°C. Store at -20°C for one year. Avoid repeated freeze/thaw cycles.		Subcellular Location: Cytoplasm ,Nucleus
Background: Molybdenum cofactor biosynthesis is a conserved pathway leading to the biological activation of molybdenum. The protein encoded by this gene is involved in this pathway. This gene was originally thought to produce a bicistronic mRNA with the potential to produce two proteins (MOCS1A and MOCS1B) from adjacent open reading frames. However, only the first open reading frame (MOCS1A) has been found to encode a protein from the putative bicistronic mRNA, whereas additional splice variants, whose full- length natures have yet to be determined, are likely to produce a fusion between the two open reading frames. This gene is defective in patients with molybdenum cofactor deficiency, type A. A related pseudogene has been identified on chromosome 16. [provided by RefSeq, Jan 2010]		e I