

www.bioss.com.cn sales@bioss.com.cn techsupport@bioss.com.cn 400-901-9800

ZO-1 Rabbit pAb

Catalog Number: bs-34023R

Target Protein: ZO-1 Concentration: 1mg/ml

> Form: Liquid Host: Rabbit

Clonality: Polyclonal

Isotype: IgG

Applications: WB (1:500-2000)

Reactivity: Human Predicted MW: 191 kDa Entrez Gene: 7082 Swiss Prot: Q07157

Purification: affinity purified by Protein A

Storage: 0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol.

Store at -20°C for one year. Avoid repeated freeze/thaw cycles. The lyophilized antibody is stable at room temperature for at least one month and for greater than a year when kept at -20°C. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is

stable for at least two weeks at 2-4°C.

Background: This gene encodes a member of the membrane-associated guanylate kinase (MAGUK)

family of proteins, and acts as a tight junction adaptor protein that also regulates


adherens junctions. Tight junctions regulate the movement of ions and

macromolecules between endothelial and epithelial cells. The multidomain structure of this scaffold protein, including a postsynaptic density 95/disc-large/zona occludens (PDZ) domain, a Src homology (SH3) domain, a guanylate kinase (GuK) domain and unique (U) motifs all help to co-ordinate binding of transmembrane proteins, cytosolic proteins, and F-actin, which are required for tight junction function. Alternative

splicing results in multiple transcript variants encoding different isoforms. [provided

by RefSeq, Aug 2017]

VALIDATION IMAGES



25 ug total protein per lane of various lysates (see on figure) probed with ZO-1 polyclonal antibody, unconjugated (bs-34023R) at 1:1000 dilution and 4°C overnight incubation. Followed by conjugated secondary antibody incubation at r.t. for 60 min.

PRODUCT SPECIFIC PUBLICATIONS

[IF=17.1] Lei Liu. et al. Myricetin Oligomer Triggers Multi-Receptor Mediated Penetration and Autophagic Restoration of Blood-Brain Barrier for Ischemic Stroke Treatment. ACS NANO. 2024;XXXX(XXX):XXX-XXX WB; MOUSE . 38533773

[IF=9.8] Yanghuan Yu. et al. MiRNA-seq and mRNA-seq revealed the mechanism of fluoride-induced cauda epididymal injury. SCI TOTAL ENVIRON. 2024 Jun;930:172895 WB,IF; Mouse . 38697545

[IF=8.8] Jiajun Chen. et al. Integrating UHPLC-MS/MS quantitative analysis and exogenous purine supplementation to elucidate the antidepressant mechanism of Chaigui granules by regulating purine metabolism. J PHARM ANAL. 2023 Aug;: WB; Rat. 10.1016/j.jpha.2023.08.008

[IF=8.039] Yifan Zhu. et al. Discovery of Selective P2Y6R Antagonists with High Affinity and In Vivo Efficacy for Inflammatory Disease Therapy. J MED CHEM. 2023;XXXX(XXX):XXX-XXX WB; Mouse . 37078976

[IF=8.2] Xinyun Qin. et al. Regulation of the intestinal flora using polysaccharides from Callicarpa nudiflora Hook to alleviate ulcerative colitis and the molecular mechanisms involved. INT J BIOL MACROMOL. 2024 Feb;258:128887 WB; MOUSE . 38118262