

bs-3386R**[Primary Antibody]****Phospho-NR1D1 (Ser55 + Ser59) Rabbit pAb****Bioss**
ANTIBODIES

www.bioss.com.cn


sales@bioss.com.cn

techsupport@bioss.com.cn


400-901-9800

— DATASHEET —**Host:** Rabbit**Isotype:** IgG**Clonality:** Polyclonal**GeneID:** 9572**Target:** Phospho-NR1D1 (Ser55 + Ser59)**Immunogen:** KLH conjugated Synthesised phosphopeptide derived from human REV-ERB alpha around the phosphorylation site of Ser55/59: PP(p-S)PTG(p-S)LT.**Purification:** affinity purified by Protein A**Concentration:** 1mg/ml**Storage:** 0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol.


Shipped at 4°C. Store at -20°C for one year. Avoid repeated freeze/thaw cycles.

Background: NR1D1, a NR1 Thyroid Hormone-Like Receptor, is encoded by the same genomic locus as, but transcribed from the opposite strand of, Thyroid Hormone Receptor Alpha (TR Alpha). NR1D1 is a target of Nuclear Receptor ROR Alpha and a transcription regulator that has been shown to affect myocyte differentiation, adipogenesis, and lipoprotein metabolism. Mice lacking NR1D1 show abnormal postnatal cerebellar development. NR1D1 expression has been documented in human skeletal muscle and a variety of mouse and rat tissues. ESTs have been isolated from human tissue libraries, including cancerous adrenal, blood, brain, breast, colon, duodenum, fetus, head/neck, kidney, lung, skeletal muscle, skin, synovium, uterus, normal brain, breast, colon, eye, heart, pancreas, pituitary, prostate, skeletal muscle, skin, testis and thyroid.**Applications:** IHC-P (1:100-500)**IHC-F** (1:100-500)**IF** (1:100-500)**Flow-Cyt** (1ug/Test)**Reactivity:** Human (predicted: Mouse, Rat, Sheep, Cow)**Predicted MW.:** 67 kDa**Subcellular Location:** Cytoplasm ,Nucleus**— VALIDATION IMAGES —**

Tissue/cell: human laryngeal tissue; 4% Paraformaldehyde-fixed and paraffin-embedded; Antigen retrieval: citrate buffer (0.01M, pH 6.0), Boiling bathing for 15min; Block endogenous peroxidase by 3% Hydrogen peroxide for 30min; Blocking buffer (normal goat serum, C-0005) at 37°C for 20 min; Incubation: Anti-Phospho-NR1D1 (Ser55 + Ser59) Polyclonal Antibody, Unconjugated(bs-3386R) 1:200, overnight at 4°C, followed by conjugation to the secondary antibody(SP-0023) and DAB(C-0010) staining



Tissue/cell: human laryngeal tissue; 4% Paraformaldehyde-fixed and paraffin-embedded; Antigen retrieval: citrate buffer (0.01M, pH 6.0), Boiling bathing for 15min; Block endogenous peroxidase by 3% Hydrogen peroxide for 30min; Blocking buffer (normal goat serum, C-0005) at 37°C for 20 min; Incubation: Anti-Phospho-NR1D1 (Ser55 + Ser59) Polyclonal Antibody, Unconjugated(bs-3386R) 1:200, overnight at 4°C, followed by conjugation to the secondary antibody(SP-0023) and DAB(C-0010) staining



Blank control (black line) :HeLa. Primary Antibody (green line): Rabbit Anti-Phospho-NR1D1 (Ser55 + Ser59) antibody (bs-3386R) Dilution:1ug/Test; Secondary Antibody (white blue line) : Goat anti-rabbit IgG-AF488 Dilution: 0.5ug/Test. Isotype control (orange line) : Normal Rabbit IgG Protocol The cells were fixed with 4% PFA (10min at room temperature)and then permeabilized with 90% ice-cold methanol for 20 min at -20°C, The cells were then incubated in 5%BSA to block non-specific protein-protein interactions for 30 min at room temperature .Cells stained with Primary Antibody for 30 min at room temperature. The secondary antibody used for 40 min at room temperature. Acquisition of 20,000 events was performed.