

**bs-2479R****[ Primary Antibody ]****CCL14/SCYL2 Rabbit pAb**

www.bioss.com.cn

sales@bioss.com.cn

techsupport@bioss.com.cn

400-901-9800

**— DATASHEET —**

<b>Host:</b> Rabbit	<b>Isotype:</b> IgG	<b>Applications:</b> <b>IHC-P</b> (1:100-500) <b>IHC-F</b> (1:100-500) <b>IF</b> (1:100-500) <b>ELISA</b> (1:5000-10000)  <b>Reactivity:</b> (predicted: Human, Rabbit)  <b>Predicted MW.:</b> 8/9 kDa  <b>Subcellular Location:</b> Secreted
<b>Clonality:</b> Polyclonal		
<b>GeneID:</b> 6358	<b>SWISS:</b> Q16627	
<b>Target:</b> CCL14/SCYL2		
<b>Immunogen:</b> KLH conjugated synthetic peptide derived from human CCL14: 20-93/93.		
<b>Purification:</b> affinity purified by Protein A		
<b>Concentration:</b> 1mg/ml		
<b>Storage:</b> 0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol. Shipped at 4°C. Store at -20°C for one year. Avoid repeated freeze/thaw cycles.		
<b>Background:</b> This gene, chemokine (C-C motif) ligand 14, is one of several CC cytokine genes clustered on 17q11.2. The CC cytokines are secreted proteins characterized by two adjacent cysteines. The cytokine encoded by this gene induces changes in intracellular calcium concentration and enzyme release in monocytes. Multiple transcript variants encoding different isoforms have been found for this gene. Read-through transcripts are also expressed that include exons from the upstream cytokine gene, chemokine (C-C motif) ligand 15, and are represented as GeneID: 348249. [provided by RefSeq].		