bs-16748R

[Primary Antibody]

KLB Rabbit pAb



www.bioss.com.cn sales@bioss.com.cn techsupport@bioss.com.cn 400-901-9800

DATASHEET		400-901-9800
Host: Rabbit	Isotype: IgG	Applications: WB (1:500-2000)
Clonality: Polyclonal	-	IHC-P (1:100-500)
GenelD: 152831	SWISS: Q86Z14	IF (1:100-500)
Target: KLB		ICC/IF (1:100-500)
Immunogen: KLH conjugated syn 451-550/1044.	nthetic peptide derived from human KLB:	
Purification: affinity purified by Protein A		Reactivity: (predicted: Human Mouse
Concentration: 1mg/ml		Rat, Sheep)
Storage: 0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol. Shipped at 4°C. Store at -20°C for one year. Avoid repeated freeze/thaw cycles.		Predicted MW.: ^{125 kDa}
Background: BetaKlotho is a 1044 amino acid single-pass type III membrane protein that plays a key role in bile acid and cholesterol metabolism by suppressing transcription of CYP7A1 (cholesterol 7- alpha-hydroxylase), the rate-limiting enzyme in bile acid synthesis. Homozygous negative betaKlotho mice showed dramatically elevated bile acid synthesis and secretion, as well as a strong upregulation of CYP7A1 and CYP8B1 and resistance to gallstone formation. FGF19 and FGF21 require direct interaction with betaKlotho for activation, intracellular signaling and gene expression modulation. Both Klotho and betaKlotho consist of two internal repeats similar to family 1 glycosidases. BetaKlotho contains two glycosyl hydrolase 1 regions, however since these regions lack essential glutamic acid residues at specific and crucial locations, the domains appear to be inactive.		e sicial subcellular Cell membrane

• [IF=6.208] Zizhen Guo. et al. & beta;-Klotho Promotes the Development of Intrauterine Adhesions via the PI3K/AKT

Signaling Pathway. INT J MOL SCI. 2022 Jan;23(19):11294 IF;Rat. 36232594

• [IF=2.923] Yamei Hu. et al. Fibroblast growth factor 21 (FGF21) promotes porcine granulosa cell estradiol production and proliferation via PI3K/AKT/mTOR signaling. THERIOGENOLOGY. 2022 Sep;: IF,WB ;Pig. 36183492