

bs-17109R**[Primary Antibody]****Bioss**
ANTIBODIES

www.bioss.com.cn


sales@bioss.com.cn

techsupport@bioss.com.cn

400-901-9800

LAT2 Rabbit pAb**DATASHEET**

Host: Rabbit	Isotype: IgG	Applications: IHC-P (1:100-500) IHC-F (1:100-500) IF (1:100-500) Reactivity: Rat (predicted: Human, Mouse, Pig, Sheep, Dog, Horse, Orangutan) Predicted MW.: 61 kDa Subcellular Location: Cytoplasm
Clonality: Polyclonal		
GeneID: 23428	SWISS: Q9UHI5	
Target: LAT2		
Immunogen: KLH conjugated synthetic peptide derived from human LAT2: 1-100/353.		
Purification: affinity purified by Protein A		
Concentration: 1mg/ml		
Storage: 0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol. Shipped at 4°C. Store at -20°C for one year. Avoid repeated freeze/thaw cycles.		
Background: L-amino acid transporter protein-2 (LAT2), a non-glycosylated membrane protein, complexes with CD98 to contribute to reabsorption of neutral amino acids in renal epithelia and blood-tissue barriers. The gene encoding LAT2 is expressed primarily in the kidney, but also to a lesser extent in placenta, brain, liver, spleen, skeletal muscle, heart, small intestine, and lung. Transfection with the antisense sequence of LAT2 suggests that LAT2 expression plays a major role in net basolateral efflux of cysteine, and points to LAT2 as a candidate gene to modulate cysteine reabsorption. In addition, the CD98/LAT2 heterodimer associates with Integrin β 1 in intestinal epithelial cells, where ligand binding to CD98 and another cell surface molecule, ICAM-1 differentially regulates LAT2 activity, suggesting a novel mechanism by which events like cell adhesion may affect amino acid transport activity.		

VALIDATION IMAGES

Paraformaldehyde-fixed, paraffin embedded (rat brain); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Antibody incubation with (LAT2) Polyclonal Antibody, Unconjugated (bs-17109R) at 1:200 overnight at 4°C, followed by operating according to SP Kit(Rabbit) (sp-0023) instructions and DAB staining.