### bs-4889R

## [ Primary Antibody ]


# EpCAM Rabbit pAb




www.bioss.com.cn sales@bioss.com.cn techsupport@bioss.com.cn 400-901-9800

– DATASHEET –		400-901-9800
Host: Rabbit	Isotype: IgG	Applications: WB (1:500-2000) Flow-Cyt (3ug/Test)
GenelD: 4072 Target: EpCAM	<b>SWISS:</b> P16422	<b>Reactivity:</b> Human, Mouse (predicted: Rat)
Immunogen: KLH conjugated synthetic peptide derived from human EpCAM: 101-200/314. < Extracellular >		
Purification: affinity purified by Protein A		Predicted MW.: <sup>35 kDa</sup>
Concentration: 1mg/ml		Subcellular
<b>Storage:</b> 0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol. Shipped at 4°C. Store at -20°C for one year. Avoid repeated freeze/thaw cycles.		Subcellular Location: Cell membrane
<b>Background:</b> This gene encodes a carcinoma-associated antigen and is a member of a family that includes at least two type I membrane proteins. This antigen is expressed on most normal epithelial cells and gastrointestinal carcinomas and functions as a homotypic calcium-independent cell adhesion molecule. The antigen is being used as a target for immunotherapy treatment of human carcinomas. Mutations in this gene result in congenital tufting enteropathy. [provided by RefSeq, Dec 2008]		cells c being

#### - VALIDATION IMAGES -



Sample: Lane 1: Large intestine (Mouse) Lysate at 40 ug Lane 2: Small intestine (Mouse) Lysate at 40 ug Primary: Anti-EpCAM (bs-4889R) at 1/1000 dilution Secondary: IRDye800CW Goat Anti-Rabbit IgG at 1/20000 dilution Predicted band size: 35 kD Observed band size: 38 kD



Blank control: A431. Primary Antibody (green line): Rabbit Anti-TEpCAM antibody (bs-4889R) Dilution: 3µg /10^6 cells; Isotype Control Antibody (orange line): Rabbit IgG . Secondary Antibody: Goat anti-rabbit IgG-AF647 Dilution: 3µg /test. Protocol The cells were incubated in 5%BSA to block non-specific protein-protein interactions for 30 min at at room temperature .Cells stained with Primary Antibody for 30 min at room temperature. The secondary antibody used for 40 min at room temperature. Acquisition of 20,000 events was performed.

### - SELECTED CITATIONS ------

• [IF=6.389] Ravindra D. Wavhale. et al. Self-Propelled Catalytically Powered Dual-Engine Magnetic Nanobots for Rapid and Highly Efficient Capture of Circulating Fetal Trophoblasts. ADV MATER INTERFACES. 2022 Jul 06 Other ;Human. 10.1002/admi.202200522