
Rabbit Anti-Human IgA, APC conjugated

Catalog Number: bs-0360R-APC

Target Protein: Rabbit Anti-Human IgA

Concentration: 1.0 mg/ml

Form: Liquid

Host: Rabbit

Clonality: Polyclonal

Isotype: IgG

Applications: IF (1:100-1000), Flow-Cyt (1:100-1000)

Excitation spectrum: 650nm

Emission spectrum: 660nm

Not yet tested in other applications.

Optimal working dilutions must be determined by the end user.

Reactivity: Human

Purification: affinity purified by Protein A

Storage: 10 mM TBS (pH=7.4) with 1% BSA, 0.03% Proclin300 and 50% glycerol.

Store at -20°C for one year. Avoid repeated freeze/thaw cycles.

Background: Monomeric IgA constitutes 5-15 % of the serum immunoglobulins whereas dimeric IgA is localized to mucosa surfaces such as saliva, gastrointestinal secretion, bronchial fluids and milk. Mucosal IgA plays a major role in host defence by neutralising infectious agents at mucosal surfaces. The production is usually local and antigen specific IgA producing B-cells can be found in regions under the lamina propria where they mature into dimeric IgA producing plasma cells. IgA deficiency is the most common immunodeficiency that may affect both serum and mucosal produced IgA.