

**bs-2519R****[ Primary Antibody ]****CD83 Rabbit pAb****Bioss**  
**ANTIBODIES**

www.bioss.com.cn


sales@bioss.com.cn

techsupport@bioss.com.cn

400-901-9800

**— DATASHEET —**

<b>Host:</b> Rabbit <b>Clonality:</b> Polyclonal <b>GeneID:</b> 9308 <b>Target:</b> CD83 <b>Immunogen:</b> KLH conjugated synthetic peptide derived from human CD83: 121-196/196. < Cytoplasmic > <b>Purification:</b> affinity purified by Protein A <b>Concentration:</b> 1mg/ml <b>Storage:</b> 0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol. Shipped at 4°C. Store at -20°C for one year. Avoid repeated freeze/thaw cycles. <b>Background:</b> The CD83 antigen is a 186 amino acid single chain glycoprotein. This molecule is a member of the immunoglobulin superfamily and is composed of an extracellular V type immunoglobulin-like single domain, a transmembrane region, and a short, 40 amino acid cytoplasmic tail. CD83 antigen undergoes extensive post translational glycosylation, as the determined Mr is twice the predicted size of the core protein. However, CD83+ cells have a unique cell surface immunophenotype that does not correlate with that of T cells, B cells, NK cells, or cells of the myelomonocytic lineage. CD83+ cells coexpress the highest levels of MHC class II molecules, when compared with other leucocyte lineages. They also coexpress T cell markers (CD2, CD5), B cell markers (CD40, CD78), myeloid cell markers (CD13, CD33, CD36) and cytokine receptors, as well as other cell surface molecules.	<b>Isotype:</b> IgG <b>SWISS:</b> Q01151 <b>Applications:</b> WB (1:500-2000) <b>Reactivity:</b> Mouse, Rat (predicted: Human) <b>Predicted MW.:</b> 21 kDa <b>Subcellular Location:</b> Cell membrane
---	--

**— VALIDATION IMAGES —**

Sample: Lane 1: Mouse Bone tissue lysates Lane  
 2: Rat Bone tissue lysates Primary: Anti-CD83  
 (bs-2519R) at 1/1000 dilution Secondary:  
 IRDye800CW Goat Anti-Rabbit IgG at 1/20000  
 dilution Predicted band size: 21 kDa Observed  
 band size: 44 kDa

**— SELECTED CITATIONS —**

- **[IF=3.272]** Miss Hongfang Meng. et al. Overexpression of hepatocyte growth factor in dental pulp stem cells ameliorates the severity of psoriasis by reducing inflammatory responses. 2021 Jun 22 IHC ;Mouse. 34155928
- **[IF=1.48]** Zhang, Suxin, et al. "Variation and significance of secretory immunoglobulin A, interleukin 6 and dendritic cells in oral cancer." Oncology Letters. IHC ;="Human". 28454394