

**bsm-52204R****[ Primary Antibody ]****BioSS**  
ANTIBODIES

www.bioss.com.cn

sales@bioss.com.cn

techsupport@bioss.com.cn


400-901-9800

**phospho-Smad2 (Ser255) Recombinant Rabbit mAb****— DATASHEET —****Host:** Rabbit**Isotype:** IgG**Clonality:** Recombinant**CloneNo.:** 8C12**GeneID:** 4087**SWISS:** Q15796**Target:** Smad2 (Ser255)**Immunogen:** KLH conjugated Synthesised phosphopeptide derived from human Smad2 around the phosphorylation site of Ser255: TL(p-S)PV.**Purification:** affinity purified by Protein A**Concentration:** 1mg/ml**Storage:** 0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol.

Shipped at 4°C. Store at -20°C for one year. Avoid repeated freeze/thaw cycles.

**Background:** The protein encoded by this gene belongs to the SMAD, a family of proteins similar to the gene products of the Drosophila gene 'mothers against decapentaplegic' (Mad) and the C. elegans gene Sma. SMAD proteins are signal transducers and transcriptional modulators that mediate multiple signaling pathways. This protein mediates the signal of the transforming growth factor (TGF)-beta, and thus regulates multiple cellular processes, such as cell proliferation, apoptosis, and differentiation. This protein is recruited to the TGF-beta receptors through its interaction with the SMAD anchor for receptor activation (SARA) protein. In response to TGF-beta signal, this protein is phosphorylated by the TGF-beta receptors. The phosphorylation induces the dissociation of this protein with SARA and the association with the family member SMAD4. The association with SMAD4 is important for the translocation of this protein into the nucleus, where it binds to target promoters and forms a transcription repressor complex with other cofactors. This protein can also be phosphorylated by activin type 1 receptor kinase, and mediates the signal from the activin. Alternatively spliced transcript variants have been observed for this gene. [provided by RefSeq, May 2012]

**Applications:** **WB** (1:500-1000)  
**IHC-P** (1:100-500)  
**IHC-F** (1:50-200)  
**IF** (1:50-200)  
**ICC/IF** (1:50-200)

**Reactivity:** Human (predicted: Mouse)**Predicted MW.:** 58 kDa**Subcellular Location:** Cytoplasm ,Nucleus**— VALIDATION IMAGES —**

Blocking buffer: 5% NFDM/TBST Primary Ab  
 dilution: 1:1000 Primary Ab incubation  
 condition: 2 hours at room temperature  
 Secondary Ab: Goat Anti-Rabbit IgG H&L (HRP)  
 Lysate: 1: (-) HeLa; 2: (+) HeLa + OA (100 mM, 60 min) + CA (100 mM, 60 min) Protein loading  
 quantity: 20 µg Exposure time: 60 s Predicted  
 MW: 52 kDa Observed MW: 58 kDa