
TREML2 Rabbit pAb

Catalog Number: bs-2737R

Target Protein: TREML2

Concentration: 1mg/ml

Form: Liquid

Host: Rabbit

Clonality: Polyclonal

Isotype: IgG

Applications: IHC-P (1:100-500), IHC-F (1:100-500), IF (1:100-500), ELISA (1:5000-10000)

Reactivity: (predicted:Human, Mouse, Rat, Pig, Sheep, Cow, Dog, Horse)

Predicted MW: 33 kDa

Entrez Gene: 79865

Swiss Prot: Q5T2D2

Source: KLH conjugated synthetic peptide derived from human TREML2: 65-150/321.

Purification: affinity purified by Protein A

Storage: 0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol.

Shipped at 4°C. Store at -20°C for one year. Avoid repeated freeze/thaw cycles.

Background: TREML2 is expressed throughout B cell development in addition to being expressed on macrophages and neutrophils and is the only TREM molecule to be found on lymphocytes. TREML2 is expressed on B lineage cells early in development, and the highest level of expression is detected on those mature peripheral B cell subpopulations that are involved in the initial humoral immune response against bacterial pathogens. TREML2 is unique in that it lacks either the conserved transmembrane lysine residue or ITAM/ITIMs within its own cytoplasmic domain. Thus, TREML2 does not exhibit any of the features associated with classical tyrosine-based signaling.

Monocytes in the bone marrow or peripheral blood do not express detectable levels of TREML2, but its expression is up-regulated in conjunction with differentiation into macrophages. TREML2 is present on neutrophils in the bone marrow as well as the periphery, and inflammatory stimuli result in a dramatic increase in the expression of TREML2 on these cells in vivo.

TREML2 is a single-pass type I membrane protein, and it contains 1 Ig-like V-type (immunoglobulin-like) domain. It is a cell surface receptor that may play a role in the innate and adaptive immune response. TREML2 is located in a gene cluster on chromosome 6 with the single Ig variable (IgV) domain activating receptors TREM1 and TREM2, but it has distinct

structural and functional properties.