

bs-1152R**[Primary Antibody]****ATP1B2 Rabbit pAb****BioSS**
ANTIBODIES

www.bioss.com.cn


sales@bioss.com.cn

techsupport@bioss.com.cn

400-901-9800

— DATASHEET —

Host: Rabbit Clonality: Polyclonal GeneID: 482 Target: ATP1B2 Immunogen: KLH conjugated synthetic peptide derived from human ATP1b2: 201-290/290. Purification: affinity purified by Protein A Concentration: 1mg/ml Storage: 0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol. Shipped at 4°C. Store at -20°C for one year. Avoid repeated freeze/thaw cycles. Background: The protein encoded by this gene belongs to the family of Na ⁺ /K ⁺ and H ⁺ /K ⁺ ATPases beta chain proteins, and to the subfamily of Na ⁺ /K ⁺ -ATPases. Na ⁺ /K ⁺ -ATPase is an integral membrane protein responsible for establishing and maintaining the electrochemical gradients of Na and K ions across the plasma membrane. These gradients are essential for osmoregulation, for sodium-coupled transport of a variety of organic and inorganic molecules, and for electrical excitability of nerve and muscle. This enzyme is composed of two subunits, a large catalytic subunit (alpha) and a smaller glycoprotein subunit (beta). The beta subunit regulates, through assembly of alpha/beta heterodimers, the number of sodium pumps transported to the plasma membrane. The glycoprotein subunit of Na ⁺ /K ⁺ -ATPase is encoded by multiple genes. This gene encodes a beta 2 subunit. [provided by RefSeq, Jul 2008]	Isotype: IgG SWISS: P14415	Applications: WB (1:200-1000) Reactivity: Mouse, Rat (predicted: Human) Predicted MW.: 33 kDa Subcellular Location: Cell membrane
---	---	---

— VALIDATION IMAGES —

Sample: Lane 1: Mouse Cerebrum tissue lysates
Lane 2: Rat Cerebrum tissue lysates Primary:
Anti- ATPase Na⁺/ K⁺ beta 2 (bs-1152R) at 1/200
dilution Secondary: IRDye800CW Goat Anti-
Rabbit IgG at 1/20000 dilution Predicted band
size: 33 kDa Observed band size: 50 kDa

— SELECTED CITATIONS —

- **[IF=41.444]** Sun, Zeguo. et al. LINE-1 promotes tumorigenicity and exacerbates tumor progression via stimulating metabolism reprogramming in non-small cell lung cancer. MOL CANCER. 2022 Dec;21(1):1-24 WB ;Human. 35842613
- **[IF=6.304]** Shiman Zuo. et al. Establishment of a novel mesenchymal stem cell-based regimen for chronic myeloid leukemia differentiation therapy. Cell Death Dis. 2021 Feb;12(2):1-15 WB ;Human. 33627636
- **[IF=4.088]** Yanhui Jia. et al. The deacetylation of Akt by SIRT1 inhibits inflammation in macrophages and protects against sepsis. EXP BIOL MED. ;(): WB ;Human,Mouse. 37211747

Important Note: This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.

- **[IF=1.689]** Wang HY et al. Occludin Endocytosis is Involved in Disruption of the Intestinal Epithelial Barrier in a Mouse Model of Alcoholic Steatohepatitis. J Dig Dis. 2019 Jul 12. WB ;Human. 31298798