
Cdc6 Rabbit pAb

Catalog Number: bs-1390R

Target Protein: Cdc6

Concentration: 1mg/ml

Form: Liquid

Host: Rabbit

Clonality: Polyclonal

Isotype: IgG

Applications: WB (1:500-2000), IHC-P (1:100-500), IHC-F (1:100-500), IF (1:100-500), Flow-Cyt (1ug/Test), ICC/IF (1:100)

Reactivity: Human, Mouse, Rat (predicted:Cow)

Predicted MW: 63 kDa

Entrez Gene: 990

Swiss Prot: Q99741

Source: KLH conjugated synthetic peptide derived from human Cdc6: 151-300/560.


Purification: affinity purified by Protein A

Storage: 0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol.

Shipped at 4°C. Store at -20°C for one year. Avoid repeated freeze/thaw cycles.


Background: Cell cycle events are regulated by the sequential activation and deactivation of cyclin depended kinase (Cdks) and by the proteolysis of cyclins. The cell division cycle (Cdc) genes are required at various points in the cell cycle. Cdc6 is the human homolog of *Saccharomyces cerevisiae* Cdc6, which is involved in the initiation of DNA replication. Cdc6 is essential for assembling the pre-replicative complex that forms at origins of DNA replication in the G1 phase of the cell cycle in budding yeast (*S. cerevisiae*). Cdc6 is also involved in checkpoint controls that ensure that DNA replication is completed before initiation of mitosis.

VALIDATION IMAGES




Sample: K562 Cell Lysate at 30 ug Primary: Anti- Cdc6 (bs-1390R) at 1/300 dilution Secondary: IRDye800CW Goat Anti-Rabbit IgG at 1/20000 dilution Predicted band size: 63 kD Observed band size: 63 kD


Tissue/cell: rat uterus tissue; 4% Paraformaldehyde-fixed and paraffin-embedded; Antigen retrieval: citrate buffer (0.01M, pH 6.0), Boiling bathing for 15min; Block endogenous peroxidase by 3% Hydrogen peroxide for 30min; Blocking buffer (normal goat serum,C-0005) at 37°C for 20 min; Incubation: Anti-Cdc6 Polyclonal Antibody, Unconjugated(bs-1390R) 1:200, overnight at 4°C, followed by conjugation to the secondary antibody(SP-0023) and DAB(C-0010) staining



Paraformaldehyde-fixed, paraffin embedded (mouse testis); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Antibody incubation with (Cdc6) Polyclonal Antibody, Unconjugated (bs-1390R) at 1:200 overnight at 4°C, followed by operating according to SP Kit(Rabbit) (sp-0023) instructionsand DAB staining.



Blank control:Hela. Primary Antibody (green line): Rabbit Anti-Cdc6 antibody (bs-1390R) Dilution: 1ug/Test; Protocol The cells were fixed with 4% PFA (10min at room temperature)and then permeabilized with 90% ice-cold methanol for 20 min at -20°C.The cells were then incubated in 5%BSA to block non-specific protein-protein interactions for 30 min at room temperature .Cells stained with Primary Antibody for 30 min at room temperature.Acquisition of 20,000 events was performed.



Hela cell; 4% Paraformaldehyde-fixed; Triton X-100 at room temperature for 20 min; Blocking buffer (normal goat serum, C-0005) at 37°C for 20 min; Antibody incubation with (Cdc6) polyclonal Antibody, Unconjugated (bs-1390R) 1:100, 90 minutes at 37°C; followed by a conjugated Goat Anti-Rabbit IgG antibody at 37°C for 90 minutes, DAPI (blue, C02-04002) was used to stain the cell nuclei.

PRODUCT SPECIFIC PUBLICATIONS

[IF=4.522] Elsaied Elnour I et al. Bta - miR - 885 promotes proliferation and inhibits differentiation of myoblasts by targeting MyoD1. J Cell Physiol. 2020 Jan 27. WB ; Bovine . 31985035

[IF=1.422] Ran M et al. miR-26a suppresses autophagy in swine Sertoli cells by targeting ULK2.Reprod Domest Anim. 2018 Aug;53(4):864-871. WB ; Pig . 29761550