

**bs-0410R****[ Primary Antibody ]**

**Bioss**  
ANTIBODIES

www.bioss.com.cn

sales@bioss.com.cn

techsupport@bioss.com.cn


400-901-9800

## Leptin receptor Rabbit pAb


### — DATASHEET —

<b>Host:</b> Rabbit	<b>Isotype:</b> IgG	<b>Applications:</b> WB (1:500-2000) <b>IHC-P</b> (1:100-500) <b>IHC-F</b> (1:100-500) <b>IF</b> (1:100-500) <b>Flow-Cyt</b> (2ug/Test)  <b>Reactivity:</b> Human, Mouse, Rat (predicted: Sheep, Horse, Goat)  <b>Predicted MW.:</b> 26 kDa  <b>Subcellular Location:</b> Secreted ,Cell membrane
<b>Clonality:</b> Polyclonal		
<b>GeneID:</b> 3953	<b>SWISS:</b> P48357	
<b>Target:</b> Leptin receptor		
<b>Immunogen:</b> KLH conjugated synthetic peptide derived from human Leptin receptor: 101-180/262.		
<b>Purification:</b> affinity purified by Protein A		
<b>Concentration:</b> 1mg/ml		
<b>Storage:</b> 0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol. Shipped at 4°C. Store at -20°C for one year. Avoid repeated freeze/thaw cycles.		
<b>Background:</b> The protein encoded by this gene belongs to the gp130 family of cytokine receptors that are known to stimulate gene transcription via activation of cytosolic STAT proteins. This protein is a receptor for leptin (an adipocyte-specific hormone that regulates body weight), and is involved in the regulation of fat metabolism, as well as in a novel hematopoietic pathway that is required for normal lymphopoiesis. Mutations in this gene have been associated with obesity and pituitary dysfunction. Alternatively spliced transcript variants encoding different isoforms have been described for this gene. It is noteworthy that this gene and LEPROT gene (GeneID:54741) share the same promoter and the first 2 exons, however, encode distinct proteins (PMID:9207021).[provided by RefSeq, Nov 2010]		


### — VALIDATION IMAGES —




Sample: Lane 1: Cerebrum (Mouse) Lysate at 40 ug  
 Lane 2: Cerebrum (Rat) Lysate at 40 ug  
 Primary: Anti-Leptin receptor (bs-0410R) at 1/1000 dilution  
 Secondary: IRDye800CW Goat Anti-Rabbit IgG at 1/20000 dilution  
 Predicted band size: 150 kD  
 Observed band size: 150 kD



Paraformaldehyde-fixed, paraffin embedded (Rat brain); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Antibody incubation with (Leptin receptor) Polyclonal Antibody, Unconjugated (bs-0410R) at 1:400 overnight at 4°C, followed by operating according to SP Kit(Rabbit) (sp-0023) instructions and DAB staining.



Tissue/cell: Mouse Embryo tissue; 4% Paraformaldehyde-fixed and paraffin-embedded; Antigen retrieval: citrate buffer (0.01M, pH 6.0), Boiling bathing for 15min; Block endogenous peroxidase by 3% Hydrogen peroxide for 30min; Blocking buffer (normal goat serum, C-0005) at 37°C for 20 min; Incubation: Anti-Leptin receptor Polyclonal Antibody, Unconjugated (bs-0410R) 1:200, overnight at 4°C, followed by conjugation to the secondary antibody (SP-0023) and DAB (C-0010) staining



Blank control: K562. Primary Antibody (green)

**Important Note:** This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.

line): Rabbit Anti-Leptin receptor antibody  
(bs-0410R) Dilution: 2µg /10<sup>6</sup> cells; Isotype  
Control Antibody (orange line): Rabbit IgG .  
Secondary Antibody : Goat anti-rabbit IgG-AF488  
Dilution: 1µg /test. Protocol The cells were  
incubated in 5%BSA to block non-specific  
protein-protein interactions for 30 min at room  
temperature .Cells stained with Primary  
Antibody for 30 min at room temperature. The  
secondary antibody used for 40 min at room  
temperature. Acquisition of 20,000 events was  
performed.

---

## — SELECTED CITATIONS —

- **[IF=8.786]** Meiyu Wu. et al. Leptin deficiency in CD8+ T cells ameliorates non-segmental vitiligo by reducing interferon-γ and Granzyme B. FRONT IMMUNOL. 2023; 14: 1158883 FCM ;Human. 37207234
- **[IF=3.69]** Meng-fan Peng. et al. Effects of total flavonoids from Eucommia ulmoides Oliv. leaves on polycystic ovary syndrome with insulin resistance model rats induced by letrozole combined with a high-fat diet. J Ethnopharmacol. 2021 Jun;273:113947 IHC ;Rat. 33617969
- **[IF=1.931]** Wu et al. Wen-dan decoction improves negative emotions in sleep-deprived rats by regulating orexin-a and leptin expression. (2014) Evid.Based.Complement.Alternat.Med. 2014:872547 ELISA,IHC ;Rat. 24864160
- **[IF=0]** Zhang, Yun, et al. "Correlation analysis between the expressions of leptin and its receptor (ObR) and clinicopathology in endometrial cancer." Cancer Biomarkers 14.5 (2014): 353-359. IHC ;="Human" . 25171477