
Leptin receptor Rabbit pAb

Catalog Number: bs-0109R

Target Protein: Leptin receptor

Concentration: 1mg/ml

Form: Liquid

Host: Rabbit

Clonality: Polyclonal

Isotype: IgG

Applications: Flow-Cyt (1µg/Test)

Reactivity: Human, Mouse, Rat (predicted:Pig, Sheep, Cow, Dog)

Predicted MW: 126 kDa

Subcellular: Secreted ,Cell membrane

Locations:

Entrez Gene: 3953

Swiss Prot: P48357

Source: KLH conjugated synthetic peptide derived from human Leptin receptor: 901-1000/1165.


Purification: affinity purified by Protein A

Storage: 0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol.


Shipped at 4°C. Store at -20°C for one year. Avoid repeated freeze/thaw cycles.

Background: The protein encoded by this gene belongs to the gp130 family of cytokine receptors that are known to stimulate gene transcription via activation of cytosolic STAT proteins. This protein is a receptor for leptin (an adipocyte-specific hormone that regulates body weight), and is involved in the regulation of fat metabolism, as well as in a novel hematopoietic pathway that is required for normal lymphopoiesis. Mutations in this gene have been associated with obesity and pituitary dysfunction. Alternatively spliced transcript variants encoding different isoforms have been described for this gene. It is noteworthy that this gene and LEPROT gene (GeneID:54741) share the same promoter and the first 2 exons, however, encode distinct proteins (PMID:9207021).[provided by RefSeq, Nov 2010]


VALIDATION IMAGES



Blank control (Black line): Raji (Black). Primary Antibody (green line): Rabbit Anti-Leptinreceptor antibody (bs-0109R) Dilution: $1\mu\text{g} / 10^6$ cells; Isotype Control Antibody (orange line): Rabbit IgG . Secondary Antibody (white blue line): Goat anti-rabbit IgG-PE Dilution: $1\mu\text{g} / \text{test}$. Protocol The cells were fixed with 70% ice-cold methanol overnight at 4°C and then permeabilized with 0.1% PBS-Tween for 20 min at room temperature. Cells stained with Primary Antibody for 30 min at room temperature. The cells were then incubated in 1 X PBS/2%BSA/10% goat serum to block non-specific protein-protein interactions followed by the antibody for 15 min at room temperature. The secondary antibody used for 40 min at room temperature. Acquisition of 20,000 events was performed.



Cell: (mo)Hepatocyte(2% BSA at 4° blocked for 30 minutes.1% Paraformaldehyde fixed for 30 minutes,0.1%TritonX-100 permeablized for 2 minutes). Concentration:1:100. Incubation: 40 minutes. Host/Blank: Hepatocytes. Flow cytometric analysis of Rabbit Anti-Leptin receptor(long)/AF488 Conjugated antibody (bs-0109R-AF488)(green) compared with control in the absence of primary antibody (blue) followed by (mo)hepatocytes.



Blank control: K562. Primary Antibody (green line): Rabbit Anti-Leptin receptor antibody (bs-0109R) Dilution: $1\mu\text{g} / 10^6$ cells; Isotype Control Antibody (orange line): Rabbit IgG . Secondary Antibody : Goat anti-rabbit IgG-AF488 Dilution: $1\mu\text{g} / \text{test}$. Protocol The cells were incubated in 5%BSA to block non-specific protein-protein interactions for 30 min at room temperature .Cells stained with Primary Antibody for 30 min at room temperature. The secondary antibody used for 40 min at room temperature. Acquisition of 20,000 events was performed.

PRODUCT SPECIFIC PUBLICATIONS

[IF=5.037] Annamária Schaffer, et al. The ontogenies of endometrial and myometrial leptin and adiponectin receptors in pregnant rats: Their putative impact on uterine contractility. Life Sci. 2022 May;297:120465 WB ; Rat . 35271883

[IF=0] Rachmawati E et al. FRUCTOSE ALTERS THE EXPRESSION OF TYPE B LEPTIN RECEPTOR IN HYPHOTALAMUS AND INTESTINUM OF Rattus Novergicus. Journal of Islamic Medicine.2018 2 (4):26-32. IHC ; Rat . 10.18860/jim.v2i2.5776