

bs-10065R

[Primary Antibody]

Netrin 4 Rabbit pAb



www.bioss.com.cn

sales@bioss.com.cn

techsupport@bioss.com.cn

400-901-9800

— DATASHEET —

| | | |
|--|----------------------|---|
| Host: Rabbit | Isotype: IgG | Applications: WB (1:500-2000) IHC-P (1:100-500) IHC-F (1:100-500) IF (1:100-500) ICC/IF (1:100-500) ELISA (1:5000-10000) Reactivity: (predicted: Human, Mouse, Rat, Rabbit, Pig, Cow, Dog, Horse) Predicted MW.: 70 kDa Subcellular Location: Secreted ,Extracellular matrix ,Cell membrane |
| Clonality: Polyclonal | | |
| GeneID: 59277 | SWISS: Q9HB63 | |
| Target: Netrin 4 | | |
| Immunogen: KLH conjugated synthetic peptide derived from human Netrin 4: 520-628/628. | | |
| Purification: affinity purified by Protein A | | |
| Concentration: 1mg/ml | | |
| Storage: 0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol. Shipped at 4°C. Store at -20°C for one year. Avoid repeated freeze/thaw cycles. | | |
| Background: May play an important role in neural, kidney and vascular development. Promotes neurite elongation from olfactory bulb explants. | | |

— SELECTED CITATIONS —

- **[IF=3.467]** Victor H. Guaiquil. et al. Expression of axon guidance ligands and their receptors in the cornea and trigeminal ganglia and their recovery after corneal epithelium injury. EXP EYE RES. Exp Eye Res. 2022 Apr;;109054 IF ;Mouse. 35427568
- **[IF=3.412]** Qinjun Chu. et al. Midazolam Attenuates Esketamine-Induced Overactive Behaviors in Mice Before the Sedation, but Not During the Recovery. FRONT VET SCI. 2022; 9: 829747 WB,IHC ;Mouse. 35478599