bs-10016R

[Primary Antibody]

phospho-ZCWCC1 (Ser739) Rabbit pAb



www.bioss.com.cn sales@bioss.com.cn techsupport@bioss.com.cn 400-901-9800



Sample: MCF-7(Human) Cell Lysate at 30 ug K562(Human) Cell Lysate at 30 ug Primary: Antiphospho-ZCWCC1 (Ser739) (bs-10016R) at 1/1000 dilution Secondary: IRDye800CW Goat Anti-Rabbit IgG at 1/20000 dilution Predicted band size: 114 kD Observed band size: 114 kD



Paraformaldehyde-fixed, paraffin embedded (rat stomach); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Antibody incubation with (phospho-ZCWCC1 (Ser739)) Polyclonal Antibody, Unconjugated (bs-10016R) at 1:200 overnight at 4°C, followed by operating according to SP Kit(Rabbit) (sp-0023) instructionsand DAB staining.



Tissue/cell: rat pancreas tissue; 4% Paraformaldehyde-fixed and paraffinembedded; Antigen retrieval: citrate buffer (0.01M, pH 6.0), Boiling bathing for 15min; Block endogenous peroxidase by 3% Hydrogen peroxide for 30min; Blocking buffer (normal goat serum,C-0005) at 37°C for 20 min; Incubation: Anti-phospho-ZCWCC1(Ser739) Polyclonal Antibody, Unconjugated(bs-10016R) 1:200, overnight at 4°C, followed by conjugation to the secondary antibody(SP-0023) and DAB(C-0010) staining



Hela cell; 4% Paraformaldehyde-fixed; Triton

X-100 at room temperature for 20 min; Blocking buffer (normal goat serum, C-0005) at 37°C for 20 min; Antibody incubation with (Phospho-ZCWCC1 (Ser739)) polyclonal Antibody, Unconjugated (bs-10016R) 1:25, 90 minutes at 37°C; followed by a conjugated Goat Anti-Rabbit IgG antibody at 37°C for 90 minutes, DAPI (blue, C02-04002) was used to stain the cell nuclei.

- SELECTED CITATIONS -

• [IF=5.17] Wang et al. PAK1-mediated MORC2 phosphorylation promotes gastric tumorigenesis. (2015) Oncotarge. 6:9877-86 WB,IHC ;Human, Mouse. 25888627