

**bs-1394R**

**[ Primary Antibody ]**

## Jab1/COPS5 Rabbit pAb

**BioSS**  
ANTIBODIES

www.bioss.com.cn

sales@bioss.com.cn


techsupport@bioss.com.cn

400-901-9800

### — DATASHEET —

<b>Host:</b> Rabbit	<b>Isotype:</b> IgG	<b>Applications:</b> WB (1:500-2000)
<b>Clonality:</b> Polyclonal		
<b>GeneID:</b> 10987	<b>SWISS:</b> Q92905	
<b>Target:</b> Jab1/COPS5		
<b>Immunogen:</b> KLH conjugated synthetic peptide derived from human Jab1: 221-334/334.		
<b>Purification:</b> affinity purified by Protein A		
<b>Concentration:</b> 1mg/ml		
<b>Storage:</b> 0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol. Shipped at 4°C. Store at -20°C for one year. Avoid repeated freeze/thaw cycles.	<b>Reactivity:</b> Human (predicted: Mouse, Rat, Pig, Cow, Dog, Horse)	
<b>Background:</b> The protein encoded by this gene is one of the eight subunits of COP9 signalosome, a highly conserved protein complex that functions as an important regulator in multiple signaling pathways. The structure and function of COP9 signalosome is similar to that of the 19S regulatory particle of 26S proteasome. COP9 signalosome has been shown to interact with SCF-type E3 ubiquitin ligases and act as a positive regulator of E3 ubiquitin ligases. This protein is reported to be involved in the degradation of cyclin-dependent kinase inhibitor CDKN1B/p27Kip1. It is also known to be an coactivator that increases the specificity of JUN/AP1 transcription factors.	<b>Predicted MW.:</b> 37 kDa	
	<b>Subcellular Location:</b> Cytoplasm ,Nucleus	

### — VALIDATION IMAGES —



Sample: 293T Cell (Human) Lysate at 30 ug  
Primary: Anti-Jab1/COPS5 (bs-1394R) at 1/300 dilution  
Secondary: IRDye800CW Goat Anti-Rabbit IgG at 1/20000 dilution  
Predicted band size: 37kD  
Observed band size: 35kD