

**bs-1041R****[ Primary Antibody ]****MEK1 + MEK2 Rabbit pAb****BioSS**  
**ANTIBODIES**

www.bioss.com.cn


sales@bioss.com.cn

techsupport@bioss.com.cn


400-901-9800

**— DATASHEET —**

<b>Host:</b> Rabbit <b>Clonality:</b> Polyclonal <b>GeneID:</b> 407835 <b>Target:</b> MEK1 + MEK2 <b>Immunogen:</b> KLH conjugated synthetic peptide derived from human MAPKK1: 301-393/393. <b>Purification:</b> affinity purified by Protein A <b>Concentration:</b> 1mg/ml <b>Storage:</b> 0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol. Shipped at 4°C. Store at -20°C for one year. Avoid repeated freeze/thaw cycles. <b>Background:</b> The protein encoded by this gene is a dual specificity protein kinase that belongs to the MAP kinase kinase family. This kinase is known to play a critical role in mitogen growth factor signal transduction. It phosphorylates and thus activates MAPK1/ERK2 and MAPK2/ERK3. The activation of this kinase itself is dependent on the Ser/Thr phosphorylation by MAP kinase kinase kinases. Mutations in this gene cause cardiofaciocutaneous syndrome (CFC syndrome), a disease characterized by heart defects, mental retardation, and distinctive facial features similar to those found in Noonan syndrome. The inhibition or degradation of this kinase is also found to be involved in the pathogenesis of Yersinia and anthrax. A pseudogene, which is located on chromosome 7, has been identified for this gene. [provided by RefSeq, Jul 2008].	<b>Isotype:</b> IgG <b>SWISS:</b> P36507 <b>Applications:</b> IHC-P (1:100-500) <b>IHC-F</b> (1:100-500) <b>IF</b> (1:100-500) <b>Reactivity:</b> Human, Mouse, Rat (predicted: Rabbit, Pig, Cow, Chicken, Dog) <b>Predicted MW.:</b> 43 kDa <b>Subcellular Location:</b> Cell membrane ,Cytoplasm
---	--

**— VALIDATION IMAGES —**

Paraformaldehyde-fixed, paraffin embedded (Rat brain); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Antibody incubation with (MEK1 + MEK2) Polyclonal Antibody, Unconjugated (bs-1041R) at 1:500 overnight at 4°C, followed by a conjugated secondary (sp-0023) for 20 minutes and DAB staining.



Tissue/cell: human gastric carcinoma; 4% Paraformaldehyde-fixed and paraffin-embedded; Antigen retrieval: citrate buffer (0.01M, pH 6.0), Boiling bathing for 15min; Block endogenous peroxidase by 3% Hydrogen peroxide for 30min; Blocking buffer (normal goat serum, C-0005) at 37°C for 20 min; Incubation: Anti-MEK1/2 (MAPKK1) Polyclonal Antibody, Unconjugated (bs-1041R) 1:200, overnight at 4°C, followed by conjugation to the secondary antibody (SP-0023) and DAB (C-0010) staining

**— SELECTED CITATIONS —**

- **[IF=5.232]** Chen F et al. Integrated Analysis of Quantitative Proteome and Transcriptional Profiles Reveals the Dynamic Function of Maternally Expressed Proteins After Parthenogenetic Activation of Buffalo Oocyte. Mol Cell Proteomics. 2018 Oct;17(10):1875-1891. WB ;Buffalo. 30002204
- **[IF=2.5]** Shi-Jia Gao. et al. Danggui Buxue decoction alleviates primary dysmenorrhea in rats by regulating the

Important Note: This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.

- MEK1/2/ERK1/2/NF- $\kappa$ B pathway. FITOTERAPIA. 2025 Jan;180:106315 WB ;Rat. 39615702
- **[IF=3.3]** Tiantian Qi. et al. XLOC\_015548 Mitigates Skeletal Muscle Atrophy via the Gadd45g/MEK/ERK Pathway and Redox Regulation. FRONT BIOSCI-LANDMRK. 2025 Apr;30(4):36233 WB ;Mouse. 40302339
  - **[IF=0]** Wang et al. Upregulation of progranulin by Helicobacter pylori in human gastric epithelial cells via p38MAPK and MEK1/2 signaling pathway: role in epithelial cell proliferation and migration. (2011) FEMS.Immunol.Med.Microbiol. 63:82-92 WB ;Human. 21707777