

bs-4288R**[Primary Antibody]**

www.bioss.com.cn


sales@bioss.com.cn

techsupport@bioss.com.cn


400-901-9800

Adenylate cyclase 1 Rabbit pAb**— DATASHEET —**

<p>Host: Rabbit</p> <p>Clonality: Polyclonal</p> <p>GeneID: 107</p> <p>Target: Adenylate cyclase 1</p> <p>Immunogen: KLH conjugated synthetic peptide derived from human Adenylate cyclase 1: 1001-1119/1119. < Cytoplasmic ></p> <p>Purification: affinity purified by Protein A</p> <p>Concentration: 1mg/ml</p> <p>Storage: 0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol. Shipped at 4°C. Store at -20°C for one year. Avoid repeated freeze/thaw cycles.</p> <p>Background: The membrane bound adenylyl cyclases (ACs) represent one of the major families of effector enzymes for G protein coupled receptors. Eight human AC isoforms (AC-1 through AC-4) have been identified up to now and genes for at least nine adenylate cyclase (AC-1-AC-9) have been cloned characterized and sequenced. This is a calmodulin-sensitive adenylyl cyclase which may be involved in regulatory processes in the central nervous system. It may play a role in neuronal plasticity, memory acquisition and learning and is expressed predominantly in the brain, retina and adrenal medulla.</p>	<p>Applications: IHC-P (1:100-500) IHC-F (1:100-500) IF (1:100-500)</p> <p>Reactivity: Human, Rat (predicted: Mouse, Pig, Cow, Dog)</p> <p>Predicted MW.: 123 kDa</p> <p>Subcellular Location: Cell membrane</p>
--	---

— VALIDATION IMAGES —

Paraformaldehyde-fixed, paraffin embedded (rat adrenal gland); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Antibody incubation with (Adenylate cyclase 1) Polyclonal Antibody, Unconjugated (bs-4288R) at 1:400 overnight at 4°C, followed by operating according to SP Kit(Rabbit) (sp-0023) instructions and DAB staining.



Paraformaldehyde-fixed, paraffin embedded (rat brain tissue); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Antibody incubation with (Adenylate cyclase 1) Polyclonal Antibody, Unconjugated (bs-4288R) at 1:400 overnight at 4°C, followed by operating according to SP Kit(Rabbit) (sp-0023) instructions and DAB staining.