

bs-6794R**[Primary Antibody]****BioSS**
ANTIBODIES

www.bioss.com.cn


sales@bioss.com.cn

techsupport@bioss.com.cn

400-901-9800

ANT-1 Rabbit pAb**DATASHEET**

Host: Rabbit	Isotype: IgG	Applications: IHC-P (1:100-500) IHC-F (1:100-500) IF (1:100-500) Reactivity: Human (predicted: Mouse, Rat, Rabbit, Pig, Sheep, Cow, Dog) Predicted MW.: 33 kDa Subcellular Location: Cell membrane ,Cytoplasm
Clonality: Polyclonal		
GeneID: 291	SWISS: P12235	
Target: ANT-1		
Immunogen: KLH conjugated synthetic peptide derived from human ANT-1: 31-130/298.		
Purification: affinity purified by Protein A		
Concentration: 1mg/ml		
Storage: 0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol. Shipped at 4°C. Store at -20°C for one year. Avoid repeated freeze/thaw cycles.		
Background: Defects in SLC25A4 are a cause of progressive external ophthalmoplegia with mitochondrial DNA deletions autosomal dominant type 2 (PEOA2) [MIM:609283]. Progressive external ophthalmoplegia is characterized by progressive weakness of ocular muscles and levator muscle of the upper eyelid. In a minority of cases, it is associated with skeletal myopathy, which causes abnormal fatigability and even permanent muscle weakness. Ragged-red fibers and atrophy are found on muscle biopsy. A large proportion of chronic ophthalmoplegias are associated with other symptoms, leading to a multisystemic pattern of this disease. Additional symptoms are variable, and may include cataracts, hearing loss, sensory axonal neuropathy, ataxia, depression, hypogonadism, and parkinsonism.		

VALIDATION IMAGES

Tissue/cell: human lung carcinoma; 4%
Paraformaldehyde-fixed and paraffin-embedded; Antigen retrieval: citrate buffer (0.01M, pH 6.0), Boiling bathing for 15min; Block endogenous peroxidase by 3% Hydrogen peroxide for 30min; Blocking buffer (normal goat serum, C-0005) at 37°C for 20 min; Incubation: Anti-ANT-1 Polyclonal Antibody, Unconjugated(bs-6794R) 1:200, overnight at 4°C, followed by conjugation to the secondary antibody(SP-0023) and DAB(C-0010) staining

SELECTED CITATIONS

- **[IF=1.7]** Jinhua Wang, et al. Indirubin induces apoptosis in ovarian cancer cells via the mitochondrial pathway. AMERICAN JOURNAL OF TRANSLATIONAL RESEARCH. 2024 Oct 15;16(10):6119-6129. Western blot ;Human. 39544767

Important Note: This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.