

bs-11603R**[Primary Antibody]**

www.bioss.com.cn

sales@bioss.com.cn

techsupport@bioss.com.cn

400-901-9800

TBR1 Rabbit pAb**— DATASHEET —**

Host: Rabbit	Isotype: IgG	Applications: WB (1:500-2000) IHC-P (1:100-500) IHC-F (1:100-500) IF (1:100-500) ICC/IF (1:100-500) ELISA (1:5000-10000) Reactivity: (predicted: Human, Mouse, Rat, Pig, Sheep, Cow, Chicken, Dog, Horse) Predicted MW.: 74 kDa Subcellular Location: Nucleus
Clonality: Polyclonal		
GeneID: 10716	SWISS: Q16650	
Target: TBR1		
Immunogen: KLH conjugated synthetic peptide derived from human TBR1: 35-130/682.		
Purification: affinity purified by Protein A		
Concentration: 1mg/ml		
Storage: 0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol. Shipped at 4°C. Store at -20°C for one year. Avoid repeated freeze/thaw cycles.		
Background: A novel murine and human gene, TBR-1, encodes a putative transcription factor related to the Brachyury (T) gene that is expressed only in postmitotic cells. T-brain-1 (TBR-1) mRNA is largely restricted to the cerebral cortex, where, during embryogenesis, it defines different regions that give rise to the paleocortex, limbic cortex and neocortex (1-3). TBR-1, Pax-6 and Emx-1 are expressed in the mouse and chicken pallium. The pallio-subpallial boundary lies at the interface between the TBR-1 and Dlx-2 expression domains. Chicken genes homologous to these mouse genes are expressed in topologically comparable patterns during development, suggesting that mouse and chicken may have similar histogenetic specification processes and field homologies (4). CASK/LIN-2, a membrane-associated guanylate kinase, is required for EGFR localization and signaling. In adult rat brain, CASK is concentrated at neuronal synapses and binds to the cell-surface proteins. CASK can interact with TBR-1, which is involved in forebrain development. CASK enters into the nucleus and binds to a specific DNA sequence (the T-element) in a complex with TBR-1. Thus, CASK acts as a coactivator of TBR-1 to induce transcription of T-element containing genes, including reelin (5).		

— SELECTED CITATIONS —

- **[IF=4]** Xiaoting Meng. et al. Electrical stimulation induced structural 3D human engineered neural tissue with well-developed neuronal network and functional connectivity. J NEURAL ENG. 2023 Jul; ICC ;Human. 37433290