

---

## EphB2 Rabbit pAb

Catalog Number: bs-0247R

Target Protein: EphB2

Concentration: 1mg/ml

Form: Liquid

Host: Rabbit

Clonality: Polyclonal

Isotype: IgG

Applications: WB (1:500-2000), IHC-P (1:100-500), IHC-F (1:100-500), IF (1:100-500)

Reactivity: Human, Mouse, Rat (predicted:Chicken, Dog)

Predicted MW: 114 kDa

Entrez Gene: 2048

Swiss Prot: P29323

Source: KLH conjugated synthetic peptide derived from human EphB2: 101-200/1055.

Purification: affinity purified by Protein A


Storage: 0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol.

Shipped at 4°C. Store at -20°C for one year. Avoid repeated freeze/thaw cycles.


**Background:** This gene encodes a member of the Eph receptor family of receptor tyrosine kinase transmembrane glycoproteins. These receptors are composed of an N-terminal glycosylated ligand-binding domain, a transmembrane region and an intracellular kinase domain. They bind ligands called ephrins and are involved in diverse cellular processes including motility, division, and differentiation. A distinguishing characteristic of Eph-ephrin signaling is that both receptors and ligands are competent to transduce a signaling cascade, resulting in bidirectional signaling. This protein belongs to a subgroup of the Eph receptors called EphB. Proteins of this subgroup are distinguished from other members of the family by sequence homology and preferential binding affinity for membrane-bound ephrin-B ligands. Allelic variants are associated with prostate and brain cancer susceptibility. Alternative splicing results in multiple transcript variants. [provided by RefSeq, May 2015]

### VALIDATION IMAGES


---



Sample: Lane 1: Testis (Mouse) Lysate at 40 ug Lane 2: Testis (Rat) Lysate at 40 ug Lane 3: Huvec (Human) Cell Lysate at 30 ug Lane 4: U251 (Human) Cell Lysate at 30 ug Lane 5: A431 (Human) Cell Lysate at 30 ug Lane 6: U2os (Human) Cell Lysate at 30 ug Lane 7: MDA-MB-231 (Human) Cell Lysate at 30 ug Lane 8: K562 (Human) Cell Lysate at 30 ug Primary: Anti-EphB2 (bs-0247R) at 1/1000 dilution Secondary: IRDye800CW Goat Anti-Rabbit IgG at 1/20000 dilution Predicted band size: 125 kD Observed band size: 120 kD



Tissue/cell: human lung carcinoma; 4% Paraformaldehyde-fixed and paraffin-embedded; Antigen retrieval: citrate buffer ( 0.01M, pH 6.0 ), Boiling bathing for 15min; Block endogenous peroxidase by 3% Hydrogen peroxide for 30min; Blocking buffer (normal goat serum,C-0005) at 37°C for 20 min; Incubation: Anti-EphB2 R Polyclonal Antibody, Unconjugated(bs-0247R) 1:500, overnight at 4°C, followed by conjugation to the secondary antibody(SP-0023) and DAB(C-0010) staining



Tissue/cell: human colon carcinoma; 4% Paraformaldehyde-fixed and paraffin-embedded; Antigen retrieval: citrate buffer ( 0.01M, pH 6.0 ), Boiling bathing for 15min; Block endogenous peroxidase by 3% Hydrogen peroxide for 30min; Blocking buffer (normal goat serum,C-0005) at 37°C for 20 min; Incubation: Anti-EphB2 R Polyclonal Antibody, Unconjugated(bs-0247R) 1:500, overnight at 4°C, followed by conjugation to the secondary antibody(SP-0023) and DAB(C-0010) staining

## PRODUCT SPECIFIC PUBLICATIONS

**[IF=2.571]** Lijing Wanget al. 1-(4-((5-chloro-4-((2-(isopropylsulfonyl)phenyl)amino)pyrimidin-2-yl)amino)-3-methoxyphenyl)-3-(2-(dimethylamino)ethyl)imidazolidin-2-one (ZX-42), a novel ALK inhibitor, induces apoptosis and protective autophagy in H2228 cells. J Pharm Pharmacol . 2020 Oct;72(10):1370-1382. WB ; Human . 32596809