
Amyloid Precursor Protein Mouse pAb

Catalog Number: bs-0112M

Target Protein: Amyloid Precursor Protein

Concentration: 1mg/ml

Form: Liquid

Host: Mouse

Clonality: Polyclonal

Isotype: IgG

Applications: WB (1:500-2000), IHC-P (1:100-500), IHC-F (1:100-500), IF (1:100-500)

Reactivity: Human, Rat, Rabbit (predicted:Pig)

Predicted MW: 72-83 kDa

Entrez Gene: 351

Swiss Prot: P05067

Source: KLH conjugated synthetic peptide derived from the middle of human Soluble APP-beta/alpha: 381-480/770.


Purification: affinity purified by Protein A

Storage: 0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol.


Shipped at 4°C. Store at -20°C for one year. Avoid repeated freeze/thaw cycles.

Background: The cerebral and vascular plaques associated with Alzheimer's disease are mainly composed of Amyloid beta peptides. beta Amyloid is derived from cleavage of the Amyloid precursor protein and varies in length from 39 to 43 amino acids. beta Amyloid [1-40], beta Amyloid [1-42], and beta Amyloid [1-43] peptides result from cleavage of Amyloid precursor protein after residues 40, 42, and 43, respectively. The cleavage takes place by gamma-secretase during the last Amyloid precursor protein processing step. beta Amyloid [1-40], beta Amyloid [1-42], and beta Amyloid [1-43] peptides are major constituents of the plaques and tangles that occur in Alzheimer's disease. beta Amyloid antibodies and peptides have been developed as tools for elucidating the biology of Alzheimer's disease.


VALIDATION IMAGES



Sample: Lane 1: Human U87MG Cell Lysates Lane 2: Human SY5Y cell lysates Lane 3: Human U251 Cell Lysates Primary: Anti-beta Amyloid 1-16 (bs-0112M) at 1/1000 dilution Secondary: IRDye800CW Goat Anti-Mouse IgG at 1/20000 dilution Predicted band size: 4.3/83 kDa Observed band size: 130 kDa



Tissue/cell: rat brain tissue; 4% Paraformaldehyde-fixed and paraffin-embedded; Antigen retrieval: citrate buffer (0.01M, pH 6.0), Boiling bathing for 15min; Block endogenous peroxidase by 3% Hydrogen peroxide for 30min; Blocking buffer (normal goat serum,C-0005) at 37°C for 20 min; Incubation: Anti-Amyloid Precursor Protein Polyclonal Antibody, Unconjugated(bs-0112M) 1:200, overnight at 4°C, followed by conjugation to the secondary antibody(SP-0024) and DAB(C-0010) staining



Paraformaldehyde-fixed, paraffin embedded (rat brain); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Antibody incubation with (Amyloid Precursor Protein) Monoclonal Antibody, Unconjugated (bs-0112M) at 1:200 overnight at 4°C, followed by operating according to SP Kit(Mouse)(sp-0024) instructionsand DAB staining.

PRODUCT SPECIFIC PUBLICATIONS

[IF=14.3] Mengni Bao. et al. PICALM Regulating the Generation of Amyloid β -Peptide to Promote Anthracycline-Induced Cardiotoxicity. ADV SCI. 2024 Jun;;2401945 IF ; Mouse . 38935046