

**bs-5724R****[ Primary Antibody ]****STK36 Rabbit pAb****BioSS**  
**ANTIBODIES**

www.bioss.com.cn


sales@bioss.com.cn

techsupport@bioss.com.cn


400-901-9800

**— DATASHEET —****Host:** Rabbit**Isotype:** IgG**Clonality:** Polyclonal**GeneID:** 27148**SWISS:** Q9NRP7**Target:** STK36**Immunogen:** KLH conjugated synthetic peptide derived from human STK36: 21-120/1315.**Purification:** affinity purified by Protein A**Concentration:** 1mg/ml**Storage:** 0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol.


Shipped at 4°C. Store at -20°C for one year. Avoid repeated freeze/thaw cycles.

**Background:** Serine/threonine protein kinase required for postnatal development, possibly by regulating the homeostasis of cerebral spinal fluid or ciliary function. Controls the activity of the transcriptional regulators GLI1, GLI2 and GLI3 by opposing the effect of SUFU and promoting their nuclear localization. GLI2 requires an additional function of STK36 to become transcriptionally active, but the enzyme does not need to possess an active kinase catalytic site for this to occur.**Applications:** IHC-P (1:100-500)**IHC-F** (1:100-500)**IF** (1:100-500)**Flow-Cyt** (1ug/Test)**Reactivity:** Human, Mouse  
(predicted: Rat, Pig, Sheep, Dog, Horse)**Predicted MW.:** 144 kDa**Subcellular Location:** Cytoplasm ,Nucleus**— VALIDATION IMAGES —**


Paraformaldehyde-fixed, paraffin embedded (mouse testis tissue); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Antibody incubation with (STK36) Polyclonal Antibody, Unconjugated (bs-5724R) at 1:400 overnight at 4°C, followed by operating according to SP Kit(Rabbit) (sp-0023) instructions and DAB staining.



Paraformaldehyde-fixed, paraffin embedded (mouse brain tissue); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Antibody incubation with (STK36) Polyclonal Antibody, Unconjugated (bs-5724R) at 1:400 overnight at 4°C, followed by operating according to SP Kit(Rabbit) (sp-0023) instructions and DAB staining.



Blank control: HeLa. Primary Antibody (green line): Rabbit Anti-STK36 antibody (bs-5724R) Dilution: 1μg /10<sup>6</sup> cells; Isotype Control Antibody (orange line): Rabbit IgG . Secondary Antibody: Goat anti-rabbit IgG-AF647 Dilution: 1μg /test. Protocol The cells were fixed with 4% PFA (10min at room temperature) and then permeabilized with 90% ice-cold methanol for 20 min at room temperature. The cells were then incubated in 5%BSA to block non-specific protein-protein interactions for 30 min at -20°C .Cells stained with Primary Antibody for 30 min at room temperature. The secondary antibody used for 40 min at room temperature. Acquisition of 20,000 events was performed.



Blank control (Black line): HeLa (Black). Primary Antibody (green line): Rabbit Anti-STK36

**Important Note:** This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.

antibody (bs-5724R) Dilution: 1µg /10<sup>6</sup> cells;  
Isotype Control Antibody (orange line): Rabbit  
IgG . Secondary Antibody (white blue line): Goat  
anti-rabbit IgG-AF647 Dilution: 1µg /test.  
Protocol The cells were fixed with 4% PFA  
(10min at room temperature)and then  
permeabilized with 90% ice-cold methanol for  
20 min at room temperature. The cells were then  
incubated in 5%BSA to block non-specific  
protein-protein interactions for 30 min at room  
temperature .Cells stained with Primary  
Antibody for 30 min at room temperature. The  
secondary antibody used for 40 min at room  
temperature. Acquisition of 20,000 events was  
performed.