

www.bioss.com.cn sales@bioss.com.cn techsupport@bioss.com.cn 400-901-9800

## **CD2AP Rabbit pAb**

Catalog Number: bs-0512R

Target Protein: CD2AP Concentration: 1mg/ml

Form: Liquid

Host: Rabbit

Clonality: Polyclonal

Isotype: IgG

Applications: WB (1:500-2000), IHC-P (1:100-500), IHC-F (1:100-500), IF (1:100-500)

Reactivity: Mouse, Rat (predicted:Human, Rabbit, Pig, Sheep, Cow, Chicken, Dog, GuineaPig, Horse)

Predicted MW: 71 kDa Entrez Gene: 23607 Swiss Prot: Q9Y5K6

Source: KLH conjugated synthetic peptide derived from human CD2AP: 561-639/639.

Purification: affinity purified by Protein A

Storage: 0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol.

Shipped at 4°C. Store at -20°C for one year. Avoid repeated freeze/thaw cycles.

Background: This gene encodes a scaffolding molecule that regulates the actin cytoskeleton. The protein


directly interacts with filamentous actin and a variety of cell membrane proteins through multiple actin binding sites, SH3 domains, and a proline-rich region containing binding sites for SH3 domains. The cytoplasmic protein localizes to membrane ruffles, lipid rafts, and the

leading edges of cells. It is implicated in dynamic actin remodeling and membrane


trafficking that occurs during receptor endocytosis and cytokinesis. Haploinsufficiency of this gene is implicated in susceptibility to glomerular disease. [provided by RefSeq, Jul

2008].


## **VALIDATION IMAGES**



Sample: Lane 1: Mouse Kidney tissue lysates Lane 2: Rat Kidney tissue lysates Primary: Anti-CD2AP (bs-0512R) at 1/1000 dilution Secondary: IRDye800CW Goat Anti-Rabbit IgG at 1/20000 dilution Predicted band size: 71 kDa Observed band size: 77 kDa



Tissue/cell: rat lung tissue; 4% Paraformaldehyde-fixed and paraffin-embedded; Antigen retrieval: citrate buffer ( 0.01M, pH 6.0 ), Boiling bathing for 15min; Block endogenous peroxidase by 3% Hydrogen peroxide for 30min; Blocking buffer (normal goat serum,C-0005) at 37°C for 20 min; Incubation: Anti-CD2AP Polyclonal Antibody, Unconjugated(bs-0512R) 1:200, overnight at 4°C, followed by conjugation to the secondary antibody(SP-0023) and DAB(C-0010) staining



Tissue/cell: rat kidney tissue; 4% Paraformaldehyde-fixed and paraffin-embedded; Antigen retrieval: citrate buffer (0.01M, pH 6.0), Boiling bathing for 15min; Block endogenous peroxidase by 3% Hydrogen peroxide for 30min; Blocking buffer (normal goat serum,C-0005) at 37°C for 20 min; Incubation: Anti-CD2AP Polyclonal Antibody, Unconjugated(bs-0512R) 1:200, overnight at 4°C, followed by conjugation to the secondary antibody(SP-0023) and DAB(C-0010) staining



Paraformaldehyde-fixed, paraffin embedded (Mouse brain); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Antibody incubation with (CD2AP) Polyclonal Antibody, Unconjugated (bs-0512R) at 1:400 overnight at 4°C, followed by operating according to SP Kit(Rabbit) (sp-0023) instructions and DAB staining.

## PRODUCT SPECIFIC PUBLICATIONS

[IF=5.195] Lifeng Wei. et al. Shenqi granule upregulates CD2AP and  $\alpha$ -actinin4 and activates autophagy through regulation of mTOR/ULK1 pathway in MPC5 cells. J ETHNOPHARMACOL. 2023 Mar;303:115942 WB; MOUSE . 36442763

[IF=3.098] Xinying Zhang. et al. Exogenous spermine attenuates diabetic kidney injury in rats by inhibiting AMPK/mTOR signaling pathway. Int J Mol Med. 2021 Mar;47(3):1-1 WB; Rat . 33537831

[IF=2.518] Derya Karabulut. et al. A different perspective on the filtration barrier after kidney stone formation: An immunohistochemical and biochemical study. 2020 Nov 05 IHC; Rat. 33155094

[IF=2.7] Zeng Xiaoshan. et al. Hypoxia-inducible factor- $1\alpha$  attenuates renal podocyte injury in male rats in a simulated high-altitude environment by upregulating Krüppel-like factor 4 expression. EXP PHYSIOL. 2024 May;: WB,IHC; Rat . 38774964

[IF=3.4] J.I. Yue. et al. Acupuncture improve proteinuria in diabetic kidney disease rats by inhibiting ferroptosis and epithelial-mesenchymal transition. HELIYON. 2024 Jul;10: IF; Rat. 39071725