## bs-9699R

## [ Primary Antibody ]

## C20orf196 Rabbit pAb



www.bioss.com.cn sales@bioss.com.cn techsupport@bioss.com.cn 400-901-9800

– DATASHEET –		400-901-	400-901-9800	
Host: Rabbit	Isotype: IgG	Applications:	<b>IHC-P</b> (1:100-500)	
Clonality: Polyclonal			<b>IHC-F</b> (1:100-500) <b>IF</b> (1:100-500)	
GenelD: 149840	<b>SWISS:</b> Q8IYI0.1		<b>II</b> (1.100-500)	
Target: C20orf196				
Immunogen: KLH conjugated syn 41-140/205.	nthetic peptide derived from human C20orf196:			
Purification: affinity purified by Protein A		Reactivity: Rat		
Concentration: 1mg/ml		neactivity	(predicted: Human	
<b>Storage:</b> 0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol. Shipped at 4°C. Store at -20°C for one year. Avoid repeated freeze/thaw cycles.		Predicted MW.:	Mouse, Dog) Predicted MW.: 23 kDa	
<b>Background:</b> Representing about 2% of human DNA, chromosome 20 consists of approximately 63 million bases and 600 genes. Chromosome 20 contains a region with numerous genes expressed in the epididymis, which are thought important for seminal production, and some viewed as potential targets for male contraception. The PRNP gene encoding the prion protein associated with spongiform encephalopathies, like Creutzfeldt-Jakob disease, is found on chromosome 20. Amyotrophic lateral sclerosis, spinal muscular atrophy, ring chromosome 20 epilepsy syndrome and Alagille syndrome are also associated with chromosome 20. The C20orf196 gene product has been provisionally designated C20orf196 pending further characterization.		1		

## - VALIDATION IMAGES



Paraformaldehyde-fixed, paraffin embedded (Rat brain); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Antibody incubation with (C20orf196) Polyclonal Antibody, Unconjugated (bs-9699R) at 1:500 overnight at 4°C, followed by a conjugated secondary (sp-0023) for 20 minutes and DAB staining.