

bs-14519R**[Primary Antibody]****phospho-eIF2B epsilon (Ser539) Rabbit pAb****BioSS**
ANTIBODIES

www.bioss.com.cn


sales@bioss.com.cn

techsupport@bioss.com.cn


400-901-9800

— DATASHEET —

Host: Rabbit	Isotype: IgG	Applications: IHC-P (1:100-500)
Clonality: Polyclonal		IHC-F (1:100-500)
GeneID: 8893	SWISS: Q13144	IF (1:100-500)
Target: eIF2B epsilon (Ser539)		Reactivity: Human
Immunogen: KLH conjugated Synthesised phosphopeptide derived from human EFHC1 around the phosphorylation site of Ser539: PD(p-S)R.		
Purification: affinity purified by Protein A		Predicted MW.: 80 kDa
Concentration: 1mg/ml		Subcellular Location: Cytoplasm ,Nucleus
Storage: 0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol. Shipped at 4°C. Store at -20°C for one year. Avoid repeated freeze/thaw cycles.		
Background: This gene encodes one of five subunits of eukaryotic translation initiation factor 2B (EIF2B), a GTP exchange factor for eukaryotic initiation factor 2 and an essential regulator for protein synthesis. Mutations in this gene and the genes encoding other EIF2B subunits have been associated with leukoencephalopathy with vanishing white matter. [provided by RefSeq, Nov 2009]		

— VALIDATION IMAGES —

Paraformaldehyde-fixed, paraffin embedded (human cervical cancer); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Antibody incubation with (p-eIF2B epsilon (Ser539)) Polyclonal Antibody, Unconjugated (bs-14519R) at 1:400 overnight at 4°C, followed by a conjugated secondary (sp-0023) for 20 minutes and DAB staining.



Paraformaldehyde-fixed, paraffin embedded (Human liver cancer); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Antibody incubation with (p-eIF2B epsilon (Ser539)) Polyclonal Antibody, Unconjugated (bs-14519R) at 1:400 overnight at 4°C, followed by operating according to SP Kit(Rabbit) (sp-0023) instructions and DAB staining.

— SELECTED CITATIONS —

- **[IF=15.304]** Yao Lei. et al. Phytochemical natural killer cells reprogram tumor microenvironment for potent immunotherapy of solid tumors. BIOMATERIALS. 2022 Jun;;121635 WB ;Mouse. 10.1016/j.biomaterials.2022.121635