

bs-17264R**[Primary Antibody]****SCAMP4 Rabbit pAb**

www.bioss.com.cn

sales@bioss.com.cn

techsupport@bioss.com.cn

400-901-9800

— DATASHEET —

Host: Rabbit	Isotype: IgG	Applications: IHC-P (1:100-500) IHC-F (1:100-500) IF (1:100-500) ICC/IF (1:100-500) ELISA (1:5000-10000)
Clonality: Polyclonal		
GeneID: 113178	SWISS: Q969E2	
Target: SCAMP4		
Immunogen: KLH conjugated synthetic peptide derived from human SCAMP4: 131-229/229.		
Purification: affinity purified by Protein A		Reactivity: (predicted: Human, Mouse, Rat, Sheep, Cow)
Concentration: 1mg/ml		
Storage: 0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol. Shipped at 4°C. Store at -20°C for one year. Avoid repeated freeze/thaw cycles.		Predicted MW.: 26 kDa
Background: Secretory carrier membrane proteins (SCAMPs) are widely distributed integral membrane proteins implicated in membrane trafficking. Most SCAMPs (e.g., SCAMP1; MIM 606911) have N-terminal cytoplasmic NPF (arg-pro-phe) repeats, 4 central transmembrane regions, and a short C-terminal cytoplasmic tail. These SCAMPs likely have a role in endocytosis that is mediated by their NPF repeats. Other SCAMPs, such as SCAMP4, lack the NPF repeats and are therefore unlikely to function in endocytosis (summary by Fernandez-Chacon and Sudhof, 2000 [PubMed 11050114]).[supplied by OMIM, Feb 2011]		Subcellular Location: Cell membrane